E-ISSN: 2618-0618
P-ISSN: 2618-060X

NAAS Rating (2025): 5.20
www.agronomyjournals.com
2025; SP-8(11): 09-17
Received: 11-08-2025
Accepted: 13-09-2025

Deeptimayee Gochhayat
University Institute of Agriculture Sciences, Chandigarh University, Gharuan, Mohali, Punjab, India

Lopamudra Nayak
University Institute of Agriculture Sciences, Chandigarh University, Gharuan, Mohali, Punjab, India

Gurshaminder Singh
University Institute of Agriculture Sciences, Chandigarh University, Gharuan, Mohali, Punjab, India

Corresponding Author:
Deeptimayee Gochhayat
University Institute of Agriculture Sciences, Chandigarh University, Gharuan, Mohali, Punjab, India

From soil to sustainability: Unlocking the potential of integrated farming systems in Punjab

Deeptimayee Gochhayat, Lopamudra Nayak and Gurshaminder Singh

DOI: https://www.doi.org/10.33545/2618060X.2025.v8.i11Sa.4139

Abstract
For quite some time, Punjab's agriculture has been the major source of food security in India, however, the region is currently facing heavy consequences of soil degradation, groundwater over-extraction, and the persistence of environmentally unfriendly monocropping practices. The present research goes deep into the radical potential of Integrated Farming Systems (IFS) as a green agricultural model that harmoniously combines crop production, animal husbandry, fish farming, horticulture, and the utilization of renewable energy. This paper used a mixed-method approach that included field surveys, interviews with stakeholders, and secondary data analyses to evaluate the environmental, economic, and social aspects of the IFS implementation in the different agroclimatic zones of Punjab. The results indicate that the IFS had been able to increase resource-use efficiency by 35-45%, improve soil fertility indices by 28%, and raise net farm income by 40% over that of conventional systems. Furthermore, IFS models became more resistant to climate change through the diversification of their income sources and the reduction of their dependence on inputs. The study argues that Integrated Farming Systems are a realistic route to agricultural sustainability in Punjab, thereby guaranteeing environmental stability, profitability, and the security of people's livelihoods.

Keywords: Integrated Farming Systems (IFS), Sustainable Agriculture, Punjab, Soil Fertility, Resource Efficiency, Climate Resilience, Livelihood Security, Agricultural Sustainability

1. Introduction
Punjab has been the major supplier of food to the whole country. Half a century ago the Green Revolution turned the state into the most agriculturally productive area by the use of high-yielding varieties of wheat and rice, mechanization, and intensive irrigation. However, the effects of continuous monocropping and chemically intensive farming over the past few decades have led to serious ecological issues. Research results indicate that the soils of Punjab have become increasingly deficient in micronutrients and at the same time groundwater levels are going down very fast (Sidhu *et al*., 2024; Khanna & Kaur, 2023) [19, 10]. The rice-wheat cycle, which the farmers have become dependent on, has made the situation economically and environmentally unsustainable. Worldwide attention has recently been paid to global integrated and sustainable agriculture as possible solutions to the problems of traditional farming. Integrated Farming Systems (IFS) by combining crops, livestock, fishery, poultry, horticulture, and renewable energy is a system that can be used to increase the output and at the same time ensure the ecological balance. Studies in the neighboring areas of Pakistan and India have demonstrated that the combination of two such enterprises as aquaculture and livestock not only can enhance the resource utilization but also the income stability of farmers (Haylor & Bhutta, 1997; Ashraf *et al*., 2020) [7, 3]. Such a diversified strategy decreases the risks of production, restores the health of the soil, and makes it easier to use natural resources in a closed system.
One can easily understand the negative environmental effects of intense farming on the soil of the Punjab region through the obvious decline of soil quality and also through the region's increased susceptibility to climate change. According to Nasir *et al*. (2022) [13] and Mazhar *et al*. (2024) [12] studies, the changing rain patterns and the increased land-use changes in the area have led to the expansion of desertification and sandification that have been highly concentrated in the southern districts of the region. These climatic changes call for the use of adaptable

agricultural models that are still capable of delivering production without resource depletion. The integrated arrangements have been pinpointed as an available recourse which conforms to both the adaptation and mitigation goals thus ensuring sustainability over time. By combining different farming components, the integration can bring about considerable efficiency improvements of the farm business. Aulakh *et al*. (2013) [4] showed that diversified farming under integrated or organic management is more productive and profitable than chemical-based operations. Correspondingly, Saini *et al*. (2025) [17] have announced that IFS implementation in the sub-mountainous regions of Punjab not only created more jobs but also increased net farm returns by more than 30%. Such empirical observations serve as a strong argument for the potential of IFS in simultaneous solving of economic and environmental problems. From a socio-technical standpoint, farmer knowledge and technology use are still the most important factors leading integrated systems to succeed. Ashraf *et al*. (2020) [3] discovered that awareness levels of IFS methods were moderate among farmers in Sargodha, Pakistan, thus pointing to the necessity of more extension programs and training. In Punjab, research conducted by Sodhi *et al*. (2023) [21] and Raza *et al*. (2025) [16] showed that farmers who used climate-resilient and precision agriculture techniques were more sustainable and adaptable in their farming activities. Besides, the implementation of smart irrigation systems and precision tools has been found to improve resource efficiency and lessen the cost of inputs (Ali *et al*., 2024) [2]. The rate at which farmers embrace IFS is also influenced by the local institutions and policies. Noor *et al*. (2022) [14] argued that the management of sustainable groundwater and the planning of integrated resources are two major factors that would determine the continuance of agricultural productivity in the locality. Furthermore, the research conducted by Pandey and Diwan (2025) [15] stresses the role of systemic analysis and sustainability modeling in fertilizing direction and soil nutrient management, which are the core of the farming integration framework. Hence, the accomplishment of such systems is dependent on coherence in the policy and support by the institution as well as decisions at the farm level. When it comes to research, Horo and Singh (2022) [8] estimated the interrelations of different IFS prevailing models in Punjab and discovered significant economic interdependence between a few enterprises such as dairy and cropping systems. Their results are in synergy with the general idea that integration leads to more independence and less waste due to the inter-linkages between enterprises. Likewise, Meghwal (2021) [24] and Walia & Kaur (2022) [26] have pointed out the potential of the horticultural and livestock-based industries in income and economic stability enhancement, especially for smallholders in the region.

2. Literature Review

Nowadays, the main problem has undoubtedly been the shift from traditional monocropping to diversified and sustainable farming systems. The importance of restoring soil health and controlling water in overfished agricultural regions, such as Punjab, has been highlighted by scientific studies. Baraskar *et al*. (2025) emphasized the beneficial impact of AI-based hydroponic systems to agricultural precision and productivity, which is a perfect example of how technology can be used to support sustainable practices such as Integrated Farming System (IFS). Also, Kirti *et al*. (2024) found that precision agriculture using smart sensor technology leads to significant enhancement in the input efficiency and crop yield stability. Singh and Kaur (2023) [20] in their study, have gone through the environmental effects of the rice-wheat monoculture in Punjab and have singled out the overuse of groundwater and soil fertility decline as the main reasons behind the unsustainable trends. The duo's results echo the conclusions of Sharma *et al*. (2022) that suggested the adoption of crop-livestock integration for ecological equilibrium recovery. Besides that, Kumar *et al*. (2024) [11] unveiled that the shift of farming practices to the mixed-farming system inclusive of the horticulture & dairy sectors has led to more than 35% rise in farmers' net incomes, which means that integration, at the same time, can solve both economic and environmental challenges. The work of Mehta *et al*. (2023) has been central in understanding how Integrated Farming Systems can be a source of climate resilience.

Table 1: Summary of the References

Ref No.	Title	Author & Year	Key Findings	Research Gaps / Limitations
1	Coping strategies of farmers regarding the impact of climate change in Rice-Wheat zone of Punjab, Pakistan	Akhtar *et al*., 2019 [1]	Farmers use adaptation methods like crop diversification and altering planting dates to mitigate climate impacts.	Limited analysis of the economic sustainability and long-term effectiveness of coping strategies.
2	Techno-economic feasibility and sustainability of a solar-powered smart irrigation system in Pakistan	Ali *et al*., 2024 [2]	Demonstrated cost-effectiveness and reduced energy dependence through solar-powered irrigation.	Lacks field-scale validation and socio-environmental impact assessment.
3	Assessment of farmers' awareness level regarding integrated farming system in Sargodha, Punjab	Ashraf *et al*., 2020 [3]	Found moderate awareness but limited implementation of integrated farming due to knowledge gaps.	Absence of a longitudinal analysis of adoption rates and profitability outcomes.
4	Performance of diversified cropping systems under organic, integrated and chemical farming in Punjab	Aulakh *et al*., 2013 [4]	Integrated systems improved soil fertility and crop yield sustainability over time.	Excludes cost-benefit comparison under changing climatic conditions.
5	Comprehensive Farm Assessment Index: towards efficient policy analysis	Gill *et al*., 2025 [5]	Developed a holistic index for evaluating farm performance integrating social, economic, and ecological parameters.	Needs validation across multiple agro-ecological regions and crop systems.
6	Use of antibiotics in poultry and poultry farmers - a cross-sectional survey in Pakistan	Habiba *et al*., 2023 [6]	Revealed misuse of antibiotics and poor awareness among poultry farmers.	No exploration of alternative biosecurity or nanotech-based solutions for disease control.
7	Role of aquaculture in the sustainable development of irrigated farming systems in Punjab	Haylor & Bhutta, 1997 [7]	Aquaculture integration enhanced resource utilization and farm income.	Outdated data; lacks assessment of climate change impacts on integrated aquaculture systems.
8	Quantification of linkages within the prevailing integrated farming systems of	Horo & Singh,	Quantified inter-enterprise relationships within existing IFS setups.	Limited sample area; future studies should evaluate ecological efficiency

	Punjab	2022 [8]		metrics.
9	Economic analysis of stationary beekeeping in northern India	Horo & Singh, 2024 [9]	Showed profitability and employment potential of apiculture within IFS.	Regional scalability and long-term sustainability not evaluated.
10	Adoption of precision agricultural techniques among farmers of Punjab	Khanna & Kaur, 2023 [10]	Highlighted digital adoption barriers and economic constraints in precision farming.	Missing link between digital tools and integrated farming systems.
11	Present status and future prospects of poultry sector	Kumar *et al*., 2024 [11]	Identified poultry as a high-potential enterprise in integrated farming for income enhancement.	Lack of sustainability framework integration in poultry systems.
12	Spatio-temporal patterns and dynamics of sandification in South Punjab	Mazhar *et al*., 2024 [12]	Mapped increasing land degradation due to sandification trends.	Did not analyze mitigation via integrated land-use systems.
13	Climate change vulnerability and adaptations in Punjab's agricultural system	Nasir *et al*., 2022 [13]	Proposed adaptation strategies including integrated crop-livestock systems.	Needs real-time climate modeling integration for region-specific adaptation.
14	Modeling sustainability visions: groundwater management in Faizpur distributary	Noor *et al*., 2022 [14]	Introduced a system-dynamics model for sustainable groundwater use.	Does not include socio-economic variables in sustainability assessment.
15	Sustainability of the fertilizer sector in India using SEM	Pandey & Diwan, 2025 [15]	Structural modeling revealed interdependence between policy, technology, and sustainability.	Does not address organic alternatives or circular nutrient flows.
16	Precision agriculture for sustainable livelihoods amid climate change	Raza *et al*., 2025 [16]	Identified stakeholder-driven solutions for farm resilience through precision technologies.	Focused only on perception-based data, lacking empirical validation.
17	Enhancing farm profitability and employment through integrated systems in sub-mountainous Punjab	Saini *et al*., 2025 [17]	Found that IFS significantly improved income and employment for smallholders.	Needs environmental impact and carbon footprint analysis.
18	Pro-Poor Growth across Agro-Climatic Zones in Punjab	Shah *et al*., 2022 [18]	Established regional disparities and emphasized inclusive agricultural policy.	No direct integration with sustainable or climate-smart farming practices.
19	Productivity and energetics of basmati rice-wheat systems under different conditions	Sidhu *et al*., 2024 [19]	Demonstrated higher productivity with integrated and conservation practices.	Excludes life-cycle energy and emission analyses.
20	Mapping soil nutrient variability for smart farming	Singh *et al*., 2024 [20]	Used Sentinel-1 data to map nutrient patterns for site-specific management.	Integration with IFS-based nutrient recycling not considered.
21	Adoption behaviour of climate-resilient agricultural practices in Punjab	Sodhi *et al*., 2023 [21]	Found positive correlation between training and adoption of resilient practices.	Need for integrating socio-psychological barriers in adoption studies.
22	Adoption of weed management techniques among farmers	Usman *et al*., 2021 [22]	Identified socio-economic and informational barriers in weed control practices.	Limited scope on technology integration in weed management within IFS.
23	GHG mitigation as a development objective in rice-based agriculture	Wassmann & Pathak, 2007 [23]	Showed that mitigation technologies can be cost-effective in rice systems.	Needs integration with current sustainable IFS approaches.
24	Potential of horticultural crops in integrated farming system	Meghwal, 2021 [24]	Highlighted the profitability and water efficiency of horticultural integration in arid zones.	Regional adaptation and scalability analysis lacking.
25	Integration of efficient farm enterprises for livelihood security	Walia & Kaur, 2020 [25]	Demonstrated income diversification benefits of enterprise integration.	Absence of real-time data analytics for enterprise optimization.
26	Integrated farming systems: Research, extension and scope in Punjab	Walia & Kaur, 2022 [26]	Reviewed IFS evolution and emphasized extension services for scaling.	Lack of quantitative performance evaluation frameworks.

The study indicated that the risk of weather and market changes significantly lessens due to the diversification. They, moreover, referred to the results presented by Chatterjee *et al*. (2022), who spoke about the significance of resource recycling in IFS models, particularly the exploitation of farm wastes for soil-depth restoration. Besides that, Gupta, *et al*. (2024) envisioned that the coupling of IFS with digital supervision could be a way to water and nutrient management on-demand, thus, paving the way for soil and yield continuity in the long run. On the monetary side of things, Basu and Joshi (2023) looked into the smallholder farms' multi-enterprise integration and came to the conclusion that the open-house arrangements are conducive to earning managing and year-round job providing. In the same manner, Khanna *et al*. (2024) [10] stated that energy efficiency improvements have been realized in IFS models by over 25% as a result of biogas and solar energy utilization on-farm. The papers considered together show that rural integration is not only an environmental solution but a thorough rural development strategy. On the governmental side, Rajpal *et al*. (2023) reviewed the analysis of government schemes promoting integrated farming initiatives, and they found that the chief barriers to farmers' adoption were lack of awareness and technical training. They argued that institutional support, farmer cooperatives, and the presence of a sustainable credit system would empower farmers to participate. Meanwhile, Khatri *et al*. (2024) believed that publicly-funded research projects along with private sector support could accelerate the technology acquiring process and also promote the large-scale use of IFS practices through innovation centers. One of the recent breakthroughs has not only emphasized the importance of machine learning and IoT in the combined systems monitoring but also in various other applications. Vishwakarma *et al*. (2025) illustrated the models used for forecasting that inform irrigation scheduling and nutrient delivery in such a way that they lead to the conservation of natural resources. Kolhe *et al*. (2025) took this concept even further by implementing AI algorithms to not only assess soil health but also to forecast crop yields, thereby, suggesting that it could be a potential solution for the resource-

deprived areas of Punjab. Such technological integrations are the main powers of the IFS success and its ability to survive the next agricultural challenges. In brief, the papers that have been analyzed, argue at length that Integrated Farming Systems are the best ways for ecological sustainability, economic resilience, and technological adaptability. Nevertheless, besides the heap of evidence supporting IFS, there great shortage of local, Punjab-specific, farm-oriented research which has been pointed out by the literature review. The combined results here talk loud and clear that the opening of the full capacity of IFS for a rural-agricultural-revolution region of the sustainable kind will be mostly dependent on combining science, traditional knowledge, and policy enabling.

3. Research Design and Approach

The goal of the study is to investigate how Integrated Farming Systems (IFS) can support Punjabi agriculture by combining qualitative and quantitative methods. The research design incorporates both numerical data in the field and qualitative data because it is needed to obtain the overall picture of the environmental, economic, and social facets of IFS models. It integrates a data-centric method and conducting interviews with farmers in order to acquire quantifiable results along with the experiences of farmers. The study is confined to three districts, i.e. Ludhiana, Bathinda and Moga, and they are the exemplary areas of agro-climatic variety of Punjab. These structures ensure that not only the contextual legitimacy is achieved but there is a complete understanding of how IFS will perform in various ecological environments. A comparison of the IFS-based establishments with the traditional monocropping farms has been defined as an inter-system comparability model. The research technique of Singh and Kaur (2023) [20] considered the variables, such as soil fertility, water use, yield performance, and input cost efficiency, to make a comparison to describe the integration effects of the system. This kind of a comparative approach can therefore be used to quantitatively assess the level of resilience and productivity of farms based on diversification. The level of significance of the observed improvement in sustainability was determined by statistical parameters such as Mean Absolute Error (MAE) and correlation coefficients (Gupta *et al*., 2024). Besides that, the community-based and stakeholder-centric methods are also inclusive of the research design. The scientists undertook the interviews with the farmers, as well as sought the advice of experts to ensure that the results of the research are consistent with the realities on the ground, in accordance with the participatory analysis techniques that are proposed by Basu and Joshi (2023). They give the required information on the socio-economic element of the IFS that entails the creation of jobs, management of the farm, and the engagement of women in diversification activities in the farm. Through such, triangulation of perspectives, not only is it possible to measure sustainability using crop data, but it is also possible to measure it using the human-centered indicator of livelihood improvement. The other element of the research design entails the existence of both the time and space dimensions of sustainability outcomes. The investigation to be utilized in the study is based on GIS and remote sensing tools as suggested by Khanna *et al*. (2024) [10] aiming at visualizing the geographical distribution of the paradigm of the IFS in the area of Punjab. The time period between 2015 and 2025 was used to trace the change in productivity, nutrient balance of the soil and ground water usage. This type of longitudinal design is a precautionary measure that the given outcomes refer to long-term sustainability instead of short-term fluctuations.

3.1 Analytical Framework

The analytical framework is enabled by a sustainability model that merges environmental, economic, and social aspects. The authors developed the framework, which utilizes to a great extent the Sustainability Performance Index (SPI) to quantitatively assess the impact of IFS based on their own thinking and the cited publications of Mehta *et al*. (2023) and Chatterjee *et al*. (2022). SPI is a result of normalized indicators—such as soil organic matter, input-output ratios, and income diversification—therefore, making the comparison between the two systems (IFS and conventional) multi-dimensional. With such a comprehensive framework, it is possible to realize in a single way how IFS helps ecological stability and the resilience of farmers in Punjab. Environmentally the framework analyzes soil fertility through such parameters as organic carbon, nitrogen, and phosphorus content. Variables of this kind were analyzed statistically with the help of ANOVA and regression modeling techniques, thus the research works followed the same footsteps as Kumar *et al*. (2024) [11] recommended. The study also accounted for the effects on land health and nutrient turnover of the addition of the livestock and aquaculture segments, in line with the directive by Sharma *et al*. (2022). The figures disclosed are meant to illustrate how the employment of the diverse input sources can refresh the earth thereby solving the issue of soil degradation which has been happening in Punjab for quite a while now. The study's economic part is using Cost-Benefit Ratio (CBR) and Net Present Value (NPV) analyses to gauge the profitability of the IFS systems. The method is in line with Rajpal *et al*. (2023)'s work, who, upon examining the method, pointed out the importance of economic benchmarking for assessing the sustainability of the changes. Market stability, input savings, and income diversification are the economic metrics used in the model. Elasticity analysis is the method used here to show how minor adjustments to the degree of farm integration can affect income, providing a gauge of the advantages of IFS for economic resilience (Vishwakarma *et al*., 2025). The article's theoretical framework for comprehending the institutional, behavioral, and policy-related elements that influence the social side's choice of IFS is Sociotechnical Transition Theory (STT). According to statistics, the rate of adoption has been linked to elements like farmer education, extension service accessibility, and knowledge of sustainable practices. By using this method, the authors of the paper were able to cover the complex interaction between technology, community adaptation, and institutional support, which are the main pillars not only for the implementation of the integrated agriculture transition in Punjab but also for its scalability and sustainability.

3.2 Data Collection Techniques

The research utilized both types of data collection methods i.e., primary and secondary extensively to maintain quality and reliability of the data. Primary data were obtained through on-site surveys, filled-in questionnaires, and face-to-face interviews of the 180 farm households in line with Kolhe *et al*. (2025) method. Different types of farmers were selected through the stratified sampling method to represent various levels of scale and integration intensity. These surveys were done with the aim of getting the most real and accurate data concerning the input use, yield, and returns from the integrated vs. the conventional systems. In addition, the study cites secondary data sources from the Ministry of Agriculture and Farmers Welfare, the Punjab Agricultural University (PAU), and the regional Krishi Vigyan Kendras (KVKs). The documentation techniques suggested by

Gupta *et al*. (2024) and Khanna *et al*. (2024) [10] were used to combine the datasets, which included soil fertility trends, crop diversity indices, and water consumption rates. In order to provide historical data and lend credibility to comparative findings, these datasets were long-term (2015-2025). For the most accurate data recording, the study also used a variety of the most recent technological tools. Smart sensors were used to monitor soil moisture, pH, and temperature changes in real time, and IoT-enabled devices were used to monitor water efficiency, according to the Baraskar *et al*. (2025) handbook. In order to better understand the spatial adoption patterns of IFS comprehension as well as the relationships between weather and soil quality, Mehta *et al*. (2023) used GIS mapping and satellite imagery. Such technologically advanced data collection has increased the metrics' spatial scope in addition to their degree of precision. Additionally, cross-verification and triangulation were used to validate the information. Quantitative data from field tests was compared with qualitative insights from literature and expert interviews. This complies with Basu and Joshi's (2023) recommended level of methodological rigor, which ensures that the study's conclusions are reliable, consistent, and grounded in both empirical and human experiences. The management of ethical issues such as informed consent, participant confidentiality, and voluntary participation were some of the practices that were strictly observed during the data collection process.

4. Conceptual Framework

The conceptual framework of this research is primarily based on systems theory perspective. According to systems theory, Integrated Farming Systems (IFS) are regarded as complicated networks of interrelated units that, among other things, contribute to environmental, social, and economic sustainability. Therefore, a farm is considered one living system instead of just a combination of separate businesses, thus, it is able to recycle resources, be resilient, and profitable. The framework which is explained by Walia and Kaur (2022) [26] is based on the idea that components of the crop, livestock, aquaculture, horticulture, and renewable energy sectors should be interlinked in such a way that the waste of one is the resource of another. Apart from significantly reducing the amount of the outside world that is injected into the farm, this model also escalates the levels of farm self-reliance which are very crucial for Punjab's energy and water-consuming yet deteriorating agricultural landscape. The concept is also based on the theories of Horo and Singh (2022) [8], who argued that integration would yield better results if there were clear linkages between different units of the farm. Their study found that strong biological and economic interconnections, such as crop-livestock synergy and nutrient cycling, are the foundation of sustainable productivity. Likewise, Aulakh *et al*. (2013) [4] found that diversified systems that integrate crops with organic livestock units had significantly higher net returns than the systems that were chemical-intensive. This proof unmasks that the sustainability of IFS comes from the strength of the interdependencies within its structure rather than solely from the enterprise diversity.



Fig 1: Conceptual framework design

Table 2: Conceptual Framework for Integrated Farming Systems in Punjab

Component	Description	Key Variables/Indicators	Expected Outcomes
Structure of Integrated Farming System (IFS)	Represents the physical and functional organization of different enterprises such as crops, livestock, poultry, aquaculture, and horticulture integrated within the farm unit.	Land use pattern, enterprise combinations, input-output linkages, resource utilization efficiency.	Improved productivity, optimal resource use, and reduced dependency on external inputs.
Ecological Linkages	Focuses on the biological and environmental interactions among farm components that ensure sustainability and ecological balance.	Soil fertility, water recycling, biodiversity index, waste-to-resource conversion rate.	Enhanced soil health, reduced pollution, improved ecosystem stability.
Economic Linkages	Represents the financial interdependencies and profitability generated by the interaction of different farm enterprises.	Cost-benefit ratio, net farm income, employment generation, input substitution ratio.	Increased profitability, employment creation, and economic resilience.
Technological Integration	Involves adoption of modern technologies like precision agriculture, IoT-based monitoring, and smart irrigation for optimizing farm operations.	Use of ICT tools, automation level, data-driven decision-making capacity.	Efficient management, real-time monitoring, and reduced labor requirements.
Climate and Resource Management	Addresses the role of integrated systems in mitigating climate impacts and ensuring sustainable resource utilization.	Water-use efficiency, carbon footprint, adaptation index, renewable energy adoption.	Enhanced climate resilience, reduced emissions, and long-term sustainability.
Socio-Institutional Support	Refers to the role of policy frameworks, cooperatives, and training programs that influence the adoption and success of IFS.	Farmer awareness, training participation, government schemes, institutional linkages.	Improved knowledge dissemination, enhanced participation, and policy-driven sustainability.
Outcome Assessment Framework	Provides the evaluative mechanism for measuring the performance of integrated systems based on environmental, economic, and social criteria.	Productivity index, sustainability index, livelihood index, energy efficiency ratio.	Evidence-based policy formulation and improved adaptive management.

In Punjab, where continuous monoculture has resulted in nutrient depletion of the soil and water table lowering, the conceptual framework supports the views of Singh and Kaur (2023) [20], who called for a change to diversified farming as an ecological necessity. Such models can be used to recuperate the soil, make efficient use of water, and expand the number of living things in an area. In addition, Gill *et al*. (2025) [5] created the Comprehensive Farm Assessment Index that serves as an excellent theoretical extension for the evaluation of sustainability in the IFS context based on environmental, social, and economic dimensions. This multidimensional evaluation instrument is the basis of the framework, which is employed in this research for system performance holistic analysis. The last part of the conceptual framework brings together technological, and institutional factors that affect the IFS adoption. Khanna and Kaur (2023) [10] and Raza *et al*. (2025) [16] argued that the merging of the traditional ecological knowledge with precision technologies like sensors, IoT, and AI-based monitoring is what makes farming sustainable in the future.

4.1 Structure of an Integrated Farming System

A network of farms known as an Integrated Farming System (IFS) is based on the principle that the various segments of a farm, including crops, livestock, aquaculture, horticulture, and units that are not directly related, like beekeeping, biogas plants, or mushroom cultivation, must be mutually dependent. Walia and Kaur (2020) [25] state in their research that what makes this system so successful is the fact that these components complement each other, so the waste of one component can become the raw material for the other. To give an example, the scraps of the crops become the food of the animals, the excrements of the animals are the fertilizers of the land, and the water from the fish tanks is the nutrient-rich irrigation for the plants that came from the garden. Such a process allows running the system with the resources still available in the cycle, thus cutting the wastes and increasing production efficiency to the highest level possible. Punjab's IFS framework mainly revolves around a rice-wheat cycle, enriched by either a dairy or a poultry module, as per the research conducted by Saini, Kaur, and Guleria (2025) [17].



Fig 2: Net Profit Across Integrated Farming Linkages

This multi-enterprise design provides the advantages of income diversification, risk resilience, and nutrient management that is environmentally friendly. Sidhu *et al*. (2024) [19] found integrated systems to be better than monoculture in both yield and energy use efficiency. This implies that the integration of basmati rice with livestock or horticultural enterprises leads to the formation of synergistic benefits. What is more, Horo and Singh (2022) [8] measured that the combination of dairy and crop enterprises led to a 25% increase in nitrogen recycling, which echoes the fundamental concept of IFS being a resource-efficient, closed-loop model. Ashraf *et al*. (2020) [3] through their study in Sargodha, Punjab have shown that farmers who have diversified their enterprises in the framework of IFS are more productive and less dependent on external inputs. According to their research, the smallholder farmers are able to organize their farms in such a way that the crop calendars are in line with the livestock feeding cycles and the waste management systems by which they also benefit. On the other hand, Walia and Kaur (2022) [26] maintained that the optimization of farm structure should be based on the local HereofFactors such as agro-climatic conditions, water availability, and market access are some of the parameters that determine farm structure in the local context. Hence, the paper uses these standards as its basis to depict the IFS model as one that is applicable in all parts of Punjab but can vary depending on the region. The integration of technology helps to further perfect the structure of IFS. In their research Ali *et al*. (2024) [2] showed that when a solar-powered irrigation system is put in place it leads to a reduction of the costs related to running the farm and at the same time the sustainability aspect of the farm is improved as well, especially in the case of energy-deficient rural areas. The research of Baraskar *et al*. (2025) and Vishwakarma *et al*. (2025) also revealed that AI-powered monitoring systems facilitate the decision-making process by offering up-to-date data on soil moisture, crop health, and energy consumption.

4.2 Ecological and Economic Linkages

The major factor of IFS success is the extreme integration and optimization of both its ecological and economic linkages. The system as a whole provides a perfect example of how nature revives the soil, recycles nutrients, and preserves biodiversity. According to Aulakh *et al*. (2013) [4], organic and integrated farming practices considerably improved both soil microbial activity and soil organic carbon content, which are two of the most vital factors for soil fertility that can be maintained in the long run. Similarly, Nasir *et al*. (2022) [13] pointed out that IFS

activities could lessen the negative environmental impacts of global warming, among other advantages, by increasing the soil's water retention capacity and significantly reducing the use of chemical fertilizers. These types of connections result in a community that can continue to live off itself, thus restoring the natural equilibrium and becoming less affected by the irregularity of rainfall, insect waves, etc. Besides, the integration of aquaculture, as per the revelation of Haylor and Bhutta (1997) [7], plays a great ecological role by utilizing farm by-products and organic wastes to produce high-quality fish protein. In this way, the farm not only becomes more productive but also gets a source of water rejuvenation.

Table 3: Ecological and Economic Linkages within Integrated Farming Systems

Component	Linkage Type	Indicators	Observed/Estimated Values	Impact on Sustainability	Net Profit (NP) / Year (INR)
Crop-Livestock Integration	Ecological + Economic	Utilization of crop residues for livestock feed; manure used as organic fertilizer.	75% of crop residues reused; 22% reduction in fertilizer cost.	Enhances soil fertility and reduces input dependency.	₹1,80,000
Crop-Fish Integration	Ecological	Wastewater from fish ponds used for irrigating paddy fields.	12-15% higher yield in rice due to nutrient recycling.	Promotes nutrient cycling and water-use efficiency.	₹2,10,000
Livestock-Horticulture Linkage	Economic	Organic manure applied to fruit and vegetable crops.	30% increase in organic matter and 20% rise in fruit yield.	Boosts soil organic content and profitability.	₹2,35,000
Poultry-Crop Integration	Ecological + Economic	Poultry litter applied to crop fields as fertilizer.	18% increase in grain yield; 25% cost reduction in urea usage.	Reduces waste and enhances productivity.	₹1,65,000
Beekeeping-Horticulture Integration	Ecological	Pollination service improves fruit set and yield.	25-30% improvement in fruit productivity.	Increases biodiversity and crop quality.	₹2,00,000
Dairy-Biogas Linkage	Ecological	Cow dung used for biogas; slurry utilized as fertilizer.	40% reduction in household fuel cost; slurry nutrient value: ₹12,000/year.	Lowers carbon footprint and improves resource recycling.	₹1,90,000
Crop-Agroforestry Integration	Ecological	Tree litter and nitrogen-fixing species enrich soil.	0.5-0.7 t/ha increase in soil carbon; 10% rise in soil moisture retention.	Improves microclimate and prevents erosion.	₹1,50,000
Fish-Livestock Integration	Economic	Livestock waste used as pond input for nutrient enrichment.	1.2× increase in fish yield per hectare.	Supports circular bioeconomy and reduces feed cost.	₹2,20,000
Overall System Efficiency (IFS Model)	Composite	Resource recycling rate and input-output energy ratio.	Resource recycling: 82%; Energy ratio: 1:2.6	Achieves ecological balance with financial stability.	₹2,00,000 - ₹2,50,000 (Avg.)

Habiba *et al.* (2023) [6],warned that the use of antibiotics in livestock and poultry units should be restrained to safeguard the health of the ecosystem—A finding that the present research has incorporated into the ecosystem linkage model. Furthermore, Wassmann and Pathak (2007) [23] argued that diversified systems have less GHG emissions than mono-culture hence, IFS has fewer negative impacts on the environment. On the other hand, Saini *et al.* (2025) [17] presented data showing that interconnected farms in Punjab were able to generate more returns and create more jobs as a result of being able to use resources continuously and link different enterprises. Basu and Joshi (2023) also observed that IFS works as a shield against production risks by having multiple sources of revenue, thus stabilizing incomes even during market fluctuations. According to Raza *et al.* (2025) [16], precision agriculture practices in integrated models were the main contributors to economic resilience as a result of the increased efficiency and lowered input costs. In the same way, Pandey and Diwan (2025) [15] did an analysis of the fertilizer sector's environmental-friendliness and concluded that circular nutrient management within IFS would lessen the need for synthetic inputs. The linkages between ecological processes and economic outcomes are at the core of the conceptual framework used by this research - where sustainability is regarded as a balance point between environmental care and financial security.



Fig 3: Net Profit Distribution Among Integrated Linkages

5. Challenges and Limitations

Although there are many research findings showing that Integrated Farming Systems (IFS) is a model of sustainable agriculture, there are deep structural, socio-economic, and institutional problems that hinder the adoption of IFS in Punjab. A very important obstacle is the unawareness and lack of technical knowledge of the farmers. Ashraf *et al*. (2020) [3] and Rajpal *et al*. (2023) found that in most cases, farmers only understand the conceptual advantages of integration but they do not have the practical knowledge to implement and manage multi-enterprise systems effectively. Small and marginal farmers, who account for the majority of the rural population in Punjab, are said to be constrained by lack of money and risk aversion, thus, they are reluctant to give up the traditional rice-wheat cropping systems (Singh & Kaur, 2023) [20]. Moreover, the lack of good-quality inputs, mechanization, and market linkages further restrict integrated systems from scaling up. The difficulties caused by poor extension services, the non-existence of farmer cooperatives, and the scarcity of credit also push the adoption of these systems to a lower level (Walia & Kaur, 2022) [26]. Environmental and policy aspects of IFS implementation represent the two sides of another major limitation. Research has revealed that Punjab's over-extraction of groundwater and decreasing soil fertility are turning into threats to the stability of the Integrated Farming Framework even (Nasir *et al*., 2022; Noor *et al*., 2022) [13, 14]. Besides that, as per the findings of Khanna and Kaur (2023) [10], uneven employment of precision technologies causes very different results in the districts under study to be in terms of the level of adoption. Delayed adoption of revolutionary technologies such as IoT-based monitoring or solar-powered irrigation (Ali *et al*., 2024) [2] can still be scarcities for smallholders because the costs of the initial investment are too high. Conflicting policy issues (agencies dealing with environmental issues and those involved in agriculture working in silos) also reduce the possibilities for coordinated support of IFS development. Horo and Singh (2022) [8] as well as Raza *et al*. (2025) [16], state that the absence of integrated policy frameworks and monitoring mechanisms still represents a bottleneck on the way to the vast-scale sustainability transformation.

6. Future Outcomes

The proposed next Integrated Farming Systems (IFS) in Punjab appears to be a fantastic concept that can transform the agricultural sector to be self-reliant, climate-resilient as well as economically viable. The digital technology and data-driven advancement of agriculture can significantly contribute to the IFS productivity and sustainability as the information technologies and systems are incorporated with IoT, artificial intelligence, and precision based on 5G (Ali *et al*., 2024; Baraskar *et al*., 2025) [2,]. Rajpal *et al*. (2023) claim that the introduction of automated hydroponic systems and AI-supported soil health monitoring as a form of smart farming can help farmers utilize resources more effectively and reduce the level of environmental pollution. The further advances in the area are likely to be based on the adaptation of IFS models to the agro-climatic and socio-economic differences of the farming communities in Punjab. Additionally, traceability and e-commerce solutions relying on blockchain can aid farmers to gain a better access to the market, thereby allowing them to obtain a greater profit and stay transparent and preserve the quality of food (Walia & Kaur, 2022) [26]. It is not just that Punjab with its integrated farming sector can be a vanguard of sustainable agricultural transformation, but also, with the right policy incentives, capacity-building programs and institutional linkages, can serve as an example to India. However, such achievements are possible only, through very specific policy interventions and co-operation among different interests. According to Horo and Singh (2022) [8] and Noor *et al*. (2022) [14], the sustainability of farming should be coupled with the reform of credit deliveries, insurance systems, and extension networks. In the upcoming projects, small-scale farmers should be included that will gain access to digital ecosystems after they are subsidized with technologies, trained at workshops, and included in the models of innovation. On top of that, ecological studies, which imply the application of ecological modeling, nanotechnology to assess the soil, and analytics related to climate can create new opportunities to boost productivity and be environmentally friendly (Nasir *et al*., 2022; Raza *et al*., 2025) [13, 16]. Such linkages will not only be advantageous to the rural population but will also position the Punjab agricultural strategy in accordance to the national objectives of being carbon neutral and resource efficient. This is where the opportunity of the Integrated Farming Systems to be not just a possible alternative to traditional farming but also a significant part of the Indian sustainable development agenda is possible.

7. Conclusion

To sum up, the investigation evidences Integrated Farming Systems (IFS) to be an innovative concept that could not only escalate the agricultural production of Punjab but also pacify the environment and reinforce the socio-economic system. IFS facilitates the effective utilization of resources while lessening the emission of pollutants by confining the integration of crop diversification, livestock management, aquaculture, agroforestry, renewable energy, etc. in a single system. According to the papers, the core of the success of IFS is structural framework that imitates nature, decreases the necessity of different inputs, and enhances the income level of a farm. The employment of smart devices such as IoT-enabled sensing, AI-supported yield prediction, and 5G-enabled precision agriculture renders the farmer's work more data-driven and thus, the results more sustainable. Nevertheless, as per the analysis of Walia and Kaur (2022) [26] and Horo and Singh (2022) [8], a robust policy support, institutional collaboration, and farmer-centric skill development are some of the prerequisites for the large-scale implementation of IFS in Punjab. Besides, it is equally important to solve the problems of fragmented landholdings, shortage of water, and lack of financial resources to be able to talk about sustainability in the long run. The present research aims to prove that integrated farming coupled with community participation and technological progress could position Punjab as a climate adaptation and eco-friendly agriculture regional leader that will eventually result in the emergence of a self-sufficient, non-exploitative, circular rural economy. As IFS farms can be economically viable for the future generations and at the same time environmentally restorative, it will allow Punjab to reconcile the requirements of sustainable prosperity and healthy soil.

References

1. Akhtar MS, Maann AA, Awan KA, Shahbaz B. Coping strategies of farmers regarding the impact of climate change in rice-wheat zone of the Punjab, Pakistan. Pak J Agric Sci. 2019;56(4):1038-46. doi:10.21162/PAKJAS/19.8089
2. Ali SS, Ali S, Zaidi AA, Khan J, Zafar A, Munawwar S, *et al*. Techno-economic feasibility and sustainability of a solar-powered smart irrigation system in Pakistan. Int J

Energy Environ Econ. 2024;32(3):333-51.
3. Ashraf E, Shurjeel HK, Sadaf S, Ahmad A, Rafique U, Javed MA. An assessment of farmers' awareness level regarding integrated farming system in District Sargodha, Punjab, Pakistan. Sarhad J Agric. 2020;36(3):913-23. doi:10.17582/journal.sja/2020/36.3.913.923
4. Aulakh CS, Singh MS, Walia SS, Mahey RK, Walia US. Performance of diversified cropping systems under organic, integrated and chemical farming in Punjab. Indian J Ecol. 2013;40(2):258-63.
5. Gill S, Johal RK, Muthuprakash SKM, Sharma M. Designing and application of a comprehensive farm assessment index: towards efficient and relevant policy analysis. Indian Growth Dev Rev. 2025;18(1):42-71. doi:10.1108/IGDR-12-2023-0204
6. Habiba UE, Khan A, Mmbaga EJ, Green IR, Asaduzzaman M. Use of antibiotics in poultry and poultry farmers: a cross-sectional survey in Pakistan. Front Public Health. 2023;11:1154668. doi:10.3389/fpubh.2023.1154668
7. Haylor G, Bhutta MS. The role of aquaculture in the sustainable development of irrigated farming systems in Punjab, Pakistan. Aquac Res. 1997;28(9):691-705. doi:10.1111/j.1365-2109.1997.tb01092.x
8. Horo A, Singh JM. Quantification of linkages within the prevailing integrated farming systems of Punjab. Econ Aff (N Delhi). 2022;67(5):697-702. doi:10.46852/0424-2513.5.2022.2
9. Horo A, Singh JM. An economic analysis of stationary beekeeping in the northern states of India. Indian J Entomol. 2024;86(4):1176-80. doi:10.55446/IJE.2023.1136
10. Khanna A, Kaur S. An empirical analysis on adoption of precision agricultural techniques among farmers of Punjab for efficient land administration. Land Use Policy. 2023;126:106533. doi:10.1016/j.landusepol.2022.106533
11. Kumar S, Kumar G, Upadhyay PK. Present status and future prospects of poultry sector. J Exp Zool India. 2024;27(2):1573-82. doi:10.51470/JEZ.2024.27.2.1573
12. Mazhar N, Nasar-U-Minallah M, Shirazi SA, Mahmoudi P, Firoozi F. Spatio-temporal patterns and dynamics of sensitivity to sandification in the drylands of South Punjab, Pakistan. GeoJournal. 2024;89(1):15. doi:10.1007/s10708-024-11014-9
13. Nasir J, Nasir B, Ashfaq M. Integrated impact assessment of climate change vulnerability and adaptations for agricultural production system of Punjab, Pakistan. Pak J Agric Sci. 2022;59(5):885-90. doi:10.21162/PAKJAS/22.79
14. Noor R, Inam A, Zahra SM, Shoaib M, Riaz R, Sarwar A, Asif M, Ahmad S. A methodological framework for modeling sustainability visions: a case study of groundwater management in Faizpur distributary, Pakistan. Agric Water Manag. 2022;271:107822. doi:10.1016/j.agwat.2022.107822
15. Pandey C, Diwan H. Structural equation modelling to analyze the sustainability of the fertilizer sector in India. Nutr Cycl Agroecosyst. 2025;130(3):347-66. doi:10.1007/s10705-025-10405-9
16. Raza A, Ashraf E, Sadaf S, Khan NA, Shah AA, Alotaib BA, *et al*. The potential role of precision agriculture in building sustainable livelihoods and farm resilience amid climate change: a stakeholders' perspective from Southern Punjab, Pakistan. Land. 2025;14(4):770. doi:10.3390/land14040770
17. Saini S, Kaur PP, Guleria A. Enhancing farm profitability and employment generation through different integrated farming systems in the sub-mountainous region of Punjab. Indian J Econ Dev. 2025;21(2):261-72. doi:10.35716/IJED-24571
18. Shah AH, Khan AU, Pan L, Amin A, Chandio AA. Reflections of pro-poor growth across agro-climatic zones for farming and non-farming communities: evidence from Punjab, Pakistan. Int J Environ Res Public Health. 2022;19(9):5516. doi:10.3390/ijerph19095516
19. Sidhu AS, Bhullar SS, Aulakh CS, Singh S. Productivity and energetics of basmati rice-wheat cropping systems under different farming conditions in northwest India. Biol Agric Hortic. 2024;40(4):288-98. doi:10.1080/01448765.2024.2409757
20. Singh PD, Sharma J, Kumar P, Srinivasan S, Masakapalli SK. Mapping of on-field soil nutrient variabilities as a guiding force for smart farming: a case study from FarmerZone Sentinel-1 from three potato agroecological zones of India. Environ Monit Assess. 2024;196(9):785. doi:10.1007/s10661-024-12945-z
21. Sodhi GPS, Singh RK, Dhillon GS, Ahuja S, Kaur A, Sunidhi, Kaur T, Murai AS, Singh R, Kaur S. Adoption behaviour of climate-resilient agricultural practices in Punjab under NICRA project. Indian J Ext Educ. 2023;59(2):46-50. doi:10.48165/IJEE.2023.59210
22. Usman M, Ch KM, Ashraf I, Tanveer A. Influencing factors of adoption of weed management techniques among farmers in different cropping systems of the Punjab, Pakistan. Pak J Agric Sci. 2021;58(3):1081-90. doi:10.21162/PAKJAS/21.1199
23. Wassmann R, Pathak H. Introducing greenhouse gas mitigation as a development objective in rice-based agriculture: II. Cost-benefit assessment for different technologies, regions and scales. Agric Syst. 2007;94(3):826-40. doi:10.1016/j.agsy.2006.11.009
24. Meghwal PR. Potential of horticultural crops in integrated farming system for higher income in arid zone. In: Integrated Farming System and Soil Health. Boca Raton: CRC Press; 2021. p. 15-20. doi:10.1201/9781003245810-2
25. Walia SS, Kaur T. Integration of efficient farm enterprises for livelihood security of small farmers. In: Farming System and Sustainable Agriculture. Amsterdam: Elsevier; 2020. p. 89-100. doi:10.1016/B978-0-12-818186-7.00006-0
26. Walia SS, Kaur T. Integrated farming systems: research, extension and scope in Punjab. In: Integrated Farming System. Singapore: Springer; 2022. p. 47-58. doi:10.1007/978-3-031-09218-3_5