



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2022; 5(2): 91-94

Received: 08-05-2022

Accepted: 10-07-2022

**Rym Jaouadi**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Mohamed Elimem**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Najla Sayari**

Department of Plant Protection, Higher  
Agronomic Institute of Chott-Mariem, University  
of Sousse, Tunisia

**Maha Kalboussi**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Chaima Lahfef**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Jamel Hammadi**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Salsabil Mhamdi**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Slim Rouz**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

**Corresponding Author:**

**Rym Jaouadi**

Research Laboratory of Agricultural Production  
Systems and Sustainable Development  
LR03AGR02, Department of Agricultural  
Production, Higher School of Agriculture of  
Mograne (ESAM), Mograne, 1121, Zaghouane,  
University of Carthage, Tunisia

## Antioxidant and insecticidal properties of *Euphorbia helioscopia* L. aqueous extracts

**Rym Jaouadi, Mohamed Elimem, Najla Sayari, Maha Kalboussi, Chaima Lahfef, Jamel Hammadi, Salsabil Mhamdi and Slim Rouz**

**DOI:** <https://doi.org/10.33545/2618060X.2022.v5.i2b.114>

### Abstract

The phytochemical composition, the antioxidant and the insecticidal activities of Tunisian *Euphorbia helioscopia* L. (Euphorbiaceae) leaves and the flower's aqueous extracts were revealed. A variation in phenolic and flavonoid contents was observed between samples. Leaves aqueous extract revealed the best phenolic contents (25.2 mg GAE/g DW). However, flower extracts were characterized by the highest flavonoid contents (8.53 mg RE/g DW). The level of antioxidant capacity estimated by free radical scavenging activity (DPPH) varied significantly among organs. Leaves aqueous extract revealed the best results ( $IC_{50}=35.11 \mu\text{g/ml}$ ). *E. helioscopia* aqueous extracts were also evaluated for their insecticidal effect against *Tribolium castaneum*, with three concentrations 10 (T1), 5 (T2), and 2.5% (T3). Probit analysis revealed that flower aqueous extracts of *E. helioscopia* exhibited the lowest  $LC_{50}$  with 2.9 mg/mL.

**Keywords:** *Euphorbia helioscopia*, aqueous extracts, antioxidant, insecticidal effect

### Introduction

There is an increasing interest in medicinal plants to screen and use in the fields of agriculture, agroalimentary and pharmacology, due to their capacity as a source of natural biologically active compounds (Jaouadi *et al.*, 2022) <sup>[1]</sup>. The use of antioxidants to prevent oxidative degradation of foods by free radicals has been widely recognized. To inhibit the oxidative chain reaction, adequate antioxidants are supplied as natural or synthetic ones (Ben El Hadj Ali *et al.*, 2015) <sup>[2]</sup>. However, synthetic antioxidants have many side effects. Hence, plant based natural antioxidants act as a good source to produce a wide range of natural antioxidants. Therefore, the development and the use of natural antioxidants obtained from plants, are desired (Asha *et al.*, 2016) <sup>[3]</sup>. Being natural products and less persistent in nature, they are also eco-friendly to surrounding flora and fauna (Sanna *et al.*, 2004; Tapondjou *et al.*, 2005; Saroukolai *et al.*, 2010; Regnault-Roger *et al.*, 2012) <sup>[4, 5, 6, 7]</sup>. Many researchers have evaluated plant extracts for the management of different insect pests (Elimem *et al.*, 2019) <sup>[8]</sup>. From this perspective, there is an increasing interest to explore plant compounds or their derivatives as botanical insecticides for protecting crops (Selin-Rani *et al.* 2016) <sup>[9]</sup>.

The genus *Euphorbia*, with more than 2000 species belongs to Euphorbiaceae family. *Euphorbia helioscopia* L. also known as sun spurge, (=Sun *Euphorbia*), is an annual plant (10 to 50 cm high) with milky latex, rising with erected reddish stem, oval alternate leaves and small yellow green flowers. It has a wide distribution in Eurasia and North Africa (Su *et al.*, 2019) <sup>[10]</sup>. *E. helioscopia* gained great interest due to the biological and medical properties of its chemical compounds. Many bioactive compounds (i.e. polyphenols, steroids, lipids, and volatile oils) have been isolated and identified. Among them, diterpenoids and flavonoids are the most prominent and abundant ones (Yang *et al.*, 2021) <sup>[11]</sup>. It was reported to be widely used in folk medicine and known for its antitumor, antiviral, antibacterial, nematocidal, antifungal and antioxidant properties (Devi and Gupta, 2000; Ramezani *et al.*, 2008; Al Younes and Abdullah, 2009; Uzair *et al.*, 2009) <sup>[12, 13, 14, 15]</sup>.

Few studies were reported on *E. helioscopia*. The aim of this work was to evaluate the total phenolic and flavonoid contents from leaves and flowers of *E. helioscopia* and to assess their antioxidant and insecticidal activities.

## Materials and Methods

### Plant material

*Euphorbia helioscopia* was collected from Bir Mcherga (Latitude: 36°26'54.32''N and Longitude: 10° 04'09.32''E, Altitude 750 m) at the flowering period. Before analyses, plant organs (leaves and flowers) were separated and air dried at room temperature for two weeks, then ground to powder before analysis.

### Preparation of the plant extracts

20 grams of each organ were ground and mixed with 200 mL of distilled water. After filtration, each extract was stored at 4 °C prior to further analysis.

### Determination of Total Phenolic and Flavonoid Contents

#### Total phenolic content

The total phenolic content was determined using the method of Chetoui *et al.* (2013) [16]. 0.5 mL of diluted sample was added to 2 mL of Folin-Ciocalteu reagent. A volume of 2.5 mL of Na<sub>2</sub>CO<sub>3</sub> (7.5%) was added, after incubation for 5 min. The absorbance at 760 nm was read, after incubation for 90 min. Total phenols were expressed as gallic acid equivalents/ g DW (mg GAE/g DW)

#### Total flavonoid content

The total flavonoid content was determined using the method of Chetoui *et al.* (2013) [16]. 1 mL of diluted extract was mixed with 1 mL of 2% AlCl<sub>3</sub>. The absorbance was measured at 430 nm, after incubation for 15min. The percentage content of flavonoids was expressed as mg rutin equivalent/g DW (mg ER/g DW).

### Antioxidant activity

The antioxidant activity was carried out using free radical scavenging activity DPPH (2, 2-diphenyl-1-picrylhydrazyl), as reported by Zaouali *et al.* (2010) [17]. 3 ml of DPPH (4\*10<sup>-5</sup> M)

was added to 1 ml of diluted extracts. The absorbance was measured at 517 nm after incubation for 30min. Trolox was used as a positive control.

### Assessment of insecticidal activity

The insecticidal efficiency of *E. helioscopia* leaves and flower extracts were evaluated against *T. castaneum*. The pest specie was extracted from the infested wheat kept at the Laboratory of Entomology at the High School of Agriculture of Mograne.

Three concentrations (10 (T1), 5 (T2), and 2.5% (T3)) of *E. helioscopia* aqueous extracts were used. Filter paper disc was placed in a Petri dish and 10 adults of *T. castaneum* were placed in each Petri dish. After 24 hours the number of dead insects was recorded after 24 hours. Water was used as a negative control. All Petri dishes were stored in a climate room at 25±1°C, 60-70% relative humidity, and a photoperiod of 16:8 (L:D) h.

Mortality rates of different treatments were estimated and corrected using Abbott's formula (Abbott, 1925) [18].

### Statistical analysis

All analyses were performed in triplicate and the results were reported as means ± standard deviation of three measurements. For each analysis, the results were compared by ANOVA followed by Duncan's multiple range test using SPSS software version 26.0 for Windows.

For the insecticidal activity, results were obtained using the Probit analysis.

## Results and discussion

### Total phenolic and flavonoid contents

In the present study, total phenolic and flavonoid contents varied significantly among plant organs (Table 1). The leaves aqueous extract exhibited the best contents of polyphenols (25.2 mg GAE/g DW). However, the highest total flavonoid content was revealed in flower aqueous extract (8.53 mg ER/g DW).

**Table 1:** Total phenolic and flavonoid contents of leaves and flowers aqueous extracts

Assays	Leaves extract	Flowers extract
<b>Total phenolic and flavonoid contents</b>		
Polyphenols (mg GAE/g DW)	25.2 <sup>b</sup> ±0.0	17.23 <sup>a</sup> ±0.3
Flavonoids (mg RE/g DW)	5.03 <sup>a</sup> ±0.11	8.53 <sup>b</sup> ±0.2
<b>Antioxidant activity</b>		
DPPH (IC <sub>50</sub> µg/ml)	35.11 <sup>a</sup> ±0.0	50.12 <sup>b</sup> ±0.3

Numbers in lines followed by the same letter are not significant at  $p > 0.05$  (Duncan's multiple range test).

As compared to previous literature data, Maoulainine *et al.* (2012) [19] reported that *E. helioscopia* flower extracts showed the best total phenolic and flavonoid contents compared to leaves and stem extracts. This discrepancy could mainly be linked to genetic factors (species, organ, phenological stage, and environmental factors) (Maoulainine *et al.*, 2012) [19]. Moreover, the accumulation of phenolic compounds depends on the processes of transport involved in the distribution of these polyphenols at the plant level and the phenological organ growth (Fico *et al.*, 2020) [20].

### Antioxidant activity

As with other biological effects, phenolic compounds were reported to display antioxidant capacity. The antioxidant capacity of *E. helioscopia* samples was evaluated by the DPPH scavenging assay (Table 1).


Our data showed that the leaf's aqueous extracts revealed the best antiradical capacity (IC<sub>50</sub> = 35.11 µg/mL), which might be due to the abundance of total phenolics in this plant organ compared to flowers. Phenolic compounds have been widely known for their significant antioxidant capacities (Swallah *et al.*,

2020) [21]. Lower activity was revealed by Maoulainine *et al.* (2012) [19] for flowers and leaves methanolic *E. helioscopia* extracts (IC<sub>50</sub> = 26.66-65.25 µg/mL), from Tunisia, and even for *E. hirta* methanolic extracts (IC<sub>50</sub> = 0.2 mg/mL) (Sharma *et al.*, 2014) [22].

### Effect of *E. helioscopia* on *Tribolium castaneum*

In order to evolve environmentally safe methods for insect control, natural bioactive compounds can be used (Jbilou *et al.*, 2006) [23]. They generate toxicity, mortality, growth inhibition and suppression of the reproductive behaviour of insects (Zettler and Arthur, 2000; Lee *et al.*, 2001; Choi *et al.*, 2006) [24, 25, 26].

*E. helioscopia* aqueous extracts were evaluated for their insecticide effect against *T. castaneum*. Results revealed that percent of mortality in control Petri dishes was very low during the six first days of observation and they ranged between 0 and 20% (Figure1). Mortalities of *T. castaneum* were observed at various exposure times and concentrations of both leaves and flowers aqueous extracts. The mortality rate ranged between 15-100% and 10-100%, for leaves and flower extracts, respectively.



**Fig 1:** Effect of leaves (A) and flowers (B) aqueous extracts on *T. castaneum* (Values followed by the same letter are not significant at  $p > 0.05$  (Duncan's multiple range test).

*T. castaneum* mortality was affected by the applied concentration of extracts, as well as the exposure time. In fact, for leaves and flower extracts, mortality rates observed in T1 reached more than 60% during the second day after treatment. However, the highest mortality rate for T2, revealed 35 and 25% for leaves and flowers, respectively, during the second day. The toxicity of aqueous extracts could be related to the expanded contact between insects and *E. helioscopia* bioactive compounds

and their increased passage through insects during the period of exposure. In line with that, Maazoun *et al.* (2017) [27] reported that plant polyphenols are toxic to insects and cause rapid death. Probit analysis revealed that flowers aqueous extracts of *E. helioscopia* exhibited the lowest  $LC_{50}$  with 2.90 mg/mL (equation of the regression line:  $Y = -2.964 + 0.911 * X$  and  $LC_{90}$  with 4.31 mg/mL (Table 2).

**Table 2:**  $LC_{50}$  (mg/ml) and  $LC_{90}$  (mg/ml) values of *E. helioscopia* aqueous extracts against *T. castaneum*.

	$LC_{50}$	$LC_{90}$	Equation of the regression line
Leaves	3.97	6.05	$Y = -2.442 + 0.616 * X$
Flowers	2.90	4.31	$Y = -2.964 + 0.911 * X$

This activity can be associated with flavonoids accumulated in flower extracts. Alonso *et al.* (2002) [28] reported that flavonoids have effects on growth reduction, pupal mass, fecundity, and increasing mortality of insects. In fact, a number of flavones have been explored as feeding deterrents against many insect pests (War *et al.*, 2012) [29]. In line with that, Selin-Rani *et al.* (2016) [9] revealed that Quercetin (flavonol), isolated from *E. hirta*, produced 90% mortality at 50 ppm.

## Conclusion

Our study on the chemical composition and biological activity of Tunisian *E. helioscopia* aqueous extracts varied significantly. The leaves aqueous extracts exhibited the highest phenolic contents and the best antiradical activity. These results may highlight the use of this specie in diverse industrial fields. In addition, *E. helioscopia* showed significant insecticidal activity against *T. castaneum*. Therefore, it is a good source for insect control. It can present a substitute for damaging chemical insecticides. However, further research on the characterization



of bioactive compounds in *E. helioscopia*, should be carried out.

### Acknowledgements

The authors thank the High School of Agriculture of Mograne (ESAM), for their financial support.

### References

- Jaouadi R, Boussaid M, Zaouali Y. Essential oil composition and antioxidant activity of five Lamiaceae species. *International Journal of Multidisciplinary Research and Growth Evaluation*. 2022;3:550-555.
- Ben El-Hadj Ali I, Chaouachi M, Bahri R, Chaieb I, Boussaida M, Harzallah-Skhiri F. Chemical composition and antioxidant, antibacterial, allelopathic and insecticidal activities of essential oil of *Thymus algeriensis* Boiss. et Reut. *Industrial Crops and Products*. 2015;77:631-639.
- Asha S, Thirunavukkarasu P, Magendira M, Sadiq AM. Antioxidant Activity of *Euphorbia hirta* Linn Leaves Extracts. *European Journal of Medicinal Plants*. 2016;14(1):1-14.
- Sanna G, Bazzoni E, Moretti M. Microencapsulated essential oils active against Indian meal moth. *Boletín de Sanidad Vegetal Plagas*. 2004;30:125-132.
- Tapondjou AL, Adher C, Fontem DA, Bouda H, Recihmuth C. Bioactivities of cymol and *Eucalyptus saligna* against *Sitophilus zeamais* Motschulsky and *Tribolium confusum*. *Journal of Stored Products Research*. 2005;41:91-102.
- Saroukolai AT, Moharramipour S, Meshkatsadat MH. Insecticidal properties of *Thymus persicus* essential oil against *Tribolium castaneum* and *Sitophilus oryzae*. *Journal of Pest Science*. 2010;83:3-8.
- Regnault-Roger C, Vincen C, Arnason JT. Essential oils in insect control: Low-risk products in a high stakes world. *Annual Review of Entomology*. 2012;57:405-424.
- Elimem M, Soltani R, Dhahr W. Evaluation of insecticidal efficiency of essential oils from *Foeniculum vulgare* var. dulce waste to control *Tribolium castaneum* Herbst 1797 (Coleoptera: Tenebrionidae) and *Trogoderma granarium* Everts 1898 (Coleoptera: Dermestidae) under laboratory conditions. *Journal of new sciences, Agriculture and Biotechnology*. 2019;65(8):4125-4135.
- Selin-Rani S, Senthil-Nathan S, Thanigaivel A, Vasanth-Srinivasan P, Edwin ES, Ponsankar A, et al. Toxicity and physiological effect of quercetin on generalist herbivore, *Spodoptera litura* Fab. and a non-target earthworm *Eisenia fetida* Savigny. *Chemosphere*. 2016;165:257-267.
- Su JC, Cheng W, Song JG, Zhong YL, Huang XJ, Jiang RW, et al. Macrocyclic diterpenoids from *Euphorbia helioscopia* and their potential anti-inflammatory activity. *Journal of Natural Products*. 2019;82(10):2818-2827.
- Yang Y, Chen X, Luan F, Wang M, Wang Z, Wang J, et al. *Euphorbia helioscopia* L.: A phytochemical and pharmacological overview. *Phytochemistry*. 2021;184:12649.
- Devi LS, Gupta P. Evaluation of some plant lattices against *Heterodera cajani* on cowpea (*Vigna sinensis*). *National Academy Science Letters*. 2000;23(5-6-):65-67.
- Ramezani M, Behravan J, Arab M, Amel Farzad S. Antiviral activity of *Euphorbia helioscopia* extracts. *Journal of Biological Sciences*. 2008;8(4):809-813.
- Al-Younis NK, Abdullah AF. Isolation and antibacterial evaluation of plant extracts from some medicinal plants in Kurdistan region. *Journal of Duhok University*. 2009;12(1):250-255.
- Uzair M, Loothar BA, Choudhary BA. Biological screening of *Euphorbia helioscopia*, L. *Pakistan Journal of Pharmaceutical Sciences*. 2009;22(2):184-186.
- Chetoui I, Messaoud C, Boussaid M, Zaouali Y. Antioxidant activity, total phenolic and flavonoid content variation among Tunisian natural populations of *Rhus tripartita* (Ucria) Grande and *Rhus pentaphylla* Desf. *Industrial Crops and Products*. 2013;51:171-177.
- Zaouali Y, Bouzaine T, Boussaid M. Essential oils composition in two *Rosmarinus officinalis* L. varieties and incidence for antimicrobial and antioxidant activities. *Food Chemistry and Toxicology*. 2010;48(11):3144-3152.
- Abbott WS. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*. 1925;18(2):265-267.
- Maoulainine BML, Jelassi A, Hassen I, Ould Boukhari, OMSA. Antioxidant proprieties of methanolic and ethanolic extracts of *Euphorbia helioscopia*, (L.) aerial parts. *International Food Research Journal*. 2012;19(3):1125-1130.
- Fico G, Bilia AR, Morelli I, Tome F. Flavonoid distribution in *Pyracantha coccinea* plants at different growth phases. *Biochemical systematics and ecology*. 2000;28:673-678.
- Swallah MS, Sun H, Affoh R, Fu H, Yu H. Antioxidant Potential Overviews of Secondary Metabolites (Polyphenols) in Fruits. *International Journal of Food Science*; c2020. Article ID 9081686.
- Sharma N, Samarakoon WK, Gyawali R, Park YH, Lee SJ, Oh SJ, et al. Evaluation of the Antioxidant, Anti-Inflammatory, and Anticancer Activities of *Euphorbia hirta* Ethanolic Extract. *Molecules*. 2014;19(9):14567-14581.
- Jbilou R, Nabili A, Sayah F. Insecticidal activity of four medicinal plant extracts against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *African Journal of Biotechnology*. 2006;5(5):936-940.
- Zettler JL, Arthur FH. Chemical control of stored product insects with fumigants and residual treatments. *Crop Protection*. 2000;19(8-10):577-582.
- Lee BH, Choi WS, Lee SE, Park BS. Fumigant toxicity of essential oils and their constituent compounds towards the rice weevil, *Sitophilus oryzae* (L.). *Crop Protection*. 2001;20(4):317-320.
- Choi WS, Park BS, Lee YH, Jang DY, Yoon HY, Lee SE. Fumigant toxicities of essential oils and monoterpenes against *Lycoriella mali* adults. *Crop Protection*. 2006;25:398-401.
- Maazoun AM, Ben Hlel T, Hamdi SH, Belhadj F, Mediouni Ben Jemâab J, Mohamed Nejib Marzouki. Screening for insecticidal potential and acetylcholinesterase activity inhibition of *Urginea maritima* bulbs extract for the control of *Sitophilus oryzae* (L). *Journal of Asia-Pacific Entomology*. 2017;20(3):752-760.
- Alonso C, Ossipova S, Ossipov V. A high concentration of glucogallin, the common precursor of hydrolysable tannins, do not deter herbivores. *Journal of Insect Behaviour*. 2002;15(5):649e657.
- War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, et al. Mechanisms of Plant Defense against Insect Herbivores. *Plant Signaling & Behavior*. 2012;7(10):1306-1320.