



E-ISSN: 2618-0618  
P-ISSN: 2618-060X  
© Agronomy  
[www.agronomyjournals.com](http://www.agronomyjournals.com)  
2021; 4(2): 97-99  
Received: 13-08-2021  
Accepted: 17-10-2021

**Gurwaan Singh**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

**Dr. SN Pandey**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

**Rahul Kumar**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

**Subhash Chandra Maurya**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

**Sunil Kumar Prajapati**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

**Corresponding Author:**  
**Gurwaan Singh**  
Chandra Shekhar Azad University  
of Agriculture and Technology  
Kanpur, Kanpur, Uttar Pradesh,  
India

## Effect of weather parameters, date of sowing on performance of wheat varieties (*Triticum aestivum* L.)

**Gurwaan Singh, Dr. SN Pandey, Rahul Kumar, Subhash Chandra Maurya and Sunil Kumar Prajapati**

### Abstract

The field experiment was conducted at Student's Instructional Farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, (UP) during *Rabi* season December 2021-April 2022. In the field environment, the growth of wheat was greatly influenced by temperature changes associated with sowing dates. The field experiment laid out in split plot design (SPD) with 4 replication and 3 different date of sowing (30<sup>th</sup> Nov, 15 Dec and 30 Dec) in main plot against 4 varieties of wheat crop (HD-2733, K-307, K-9107 and HD-2967) in sub plot. Growth parameters were higher in early sown crops as compared with late sown wheat crops. Among varieties HD-2967 sown on 30<sup>th</sup> Nov obtain highest growth parameters as compare to HD-2733, K-9107 and K-307. It may be concluded that optimum sowing time of wheat crop is 30<sup>th</sup> Nov as compare to late sown wheat. Among the most promising variety.

The sowing on 30<sup>th</sup> November recorded better growth, yield attributes and grain yield (44.94 q ha<sup>-1</sup>) and harvest index (39.05%) of wheat crop. Among wheat varieties tested HD-2967 exhibited maximum grain yield (42.30 q ha<sup>-1</sup>) and harvest index (39.34 %) compared to other varieties.

**Keywords:** Productivity, sowing dates, varieties, wheat

### Introduction

Wheat (*Triticum aestivum* L.) is known as the "king of cereals" and one of the world's most significant staple food crops, being grown in at least 43 nations. Wheat provides sustenance to around 35 percent of the world's population, and it accounts for about 20% of the world's protein supply. Wheat kernels contain approximately 12 percent water, with carbohydrates (65-80 percent) primarily as starch, protein (8-15 percent) containing adequate amounts of all essential amino acids (except lysine, tryptophan, and methionine), fats (1.5- 2 percent), minerals (1.5-2 percent), vitamins (such as B complex and vitamin E), and crude fibres (2.2 percent). Wheat straw is used for cattle bedding and feed.

Other industrial uses of wheat grain include starch for paste, alcohol oil, and glut in straw. The main wheat-growing area is in India's northern region. During the growing season of the wheat crop, the optimum temperature regime is 20-22 °C at sowing, 16-22 °C at tillering until grain filling, and gradually climbs to 40 °C at harvesting (Sharma, 2000). Sowing takes place between the first week of October and the latter week of December. Delayed sowing exposes crops to adverse conditions such as low temperatures during vegetative growth, resulting in a low germination rate, poor tillering ability, and low plant population. Delayed sowing exposes the crop to adverse conditions such as low temperature during vegetative growth, resulting in a low germination rate, poor tillering ability and low plant population. The wheat sown on the normal date of sowing (25<sup>th</sup> November) attained 15 days late physiological maturity as compared to late sown condition. Delayed sowing also increases the possibility of crops exposure to high temperature during the grain filling stage, which is harmful to leaf photosynthesis, grain filling, and final yield formation. Early sowing enhanced germination unit area<sup>-1</sup>, plant height, spikelets spike<sup>-1</sup>, grains spike<sup>-1</sup> and 1000- grain weight over late sowing. The effect of date of sowing on the growth of wheat was assessed on a stand counts per m<sup>-2</sup>, plant height at various stages (cm), leaf area index at different stages, dry matter accumulation and number of tillers m<sup>-2</sup> and yield attributes viz. test weight, number of grains ear head<sup>-1</sup>, ear head m<sup>-2</sup>, length of the ear (cm). Fatima *et al.*, reported that number of different adaptation strategies based on early sowing,

irrigation and a combination of sowing and irrigation adaptations were examined to recover the potential losses that would occur due to climate change. To feed the ever-growing population, the country would need to produce 101.7 million tonnes by 2025 (Ganesh *et al.*, 2012). Under changing climate temperature stress, which is one of the most serious environmental hazards to agriculture, it is a huge issue. According to the 3rd Advance Estimates for 2020-21, overall food grain output in the country would reach a new high of 308.65 million tonnes. Food grain production is 29.77 million tonnes more than the average food grain production of the last five years. Wheat production is expected to reach a new high of 109.52 million tonnes in 2021- 22. It is higher by 9.10 million tonnes than the average wheat production of 100.42 million in 2020-21. Department of Agriculture, Cooperation, and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, 4rd Advance Estimates 2020-21).

### Methods and Materials

An experiment was carried out during the year 2021-22 (November-April) on Students' Instructional Farm (SIF) at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, (UP) Geo-graphically, Kanpur is situated in the central part of Uttar Pradesh and the sub-tropical semi-arid tract of North India. The Kanpur-city district lies between 26° 0' 29' 35'' North latitude and 80° 18' 25'' East longitude and is located on an elevation of about 125.9 meters above mean sea level in Gangetic plain. The weekly maximum and minimum temperature during experiment were 41.8°C and 4.9 °C, respectively. The mean annual rainfall is about 1194 mm. The soil was found clay loam having slightly alkaline pH and Organic carbon with a value of 0.49.

The experiment was setup according to the split-plot design with four replication, Three main plot treatments (30<sup>th</sup> November, 15<sup>th</sup> December and 30<sup>th</sup> December and four sub plot [HD-2733 (V<sub>1</sub>), [K-307 (V<sub>2</sub>), [K-9107 (V<sub>3</sub>) and [HD- 2967 (V<sub>4</sub>). Five irrigation including pre- sown were applied to the experiments. 48 plots were made for the experiments each with a gross size of 8m x 3 m = 24.0 m.

The row to row distance of 22.5 cm. To achieve accurate precision, the appropriate sampling techniques were implied for the proper balance in sampling at minimum cost. For biometric observation, plants were selected randomly in each plot to assess the possible relationship between various growth attributes and variability in climate. The observation were recorded at 30, 45, 60, 90, 105 and maturity in the marked area of plots. The land was prepared as per the requirement of the experiment.

### Result and Discussion

#### Effect of treatments on growth characters

The results revealed that growth characters *viz.* plant height, number of tillers, fresh weight, and dry weight of plant illustrate that growth of wheat varieties was significantly influenced by different dates of sowing and varieties. 30<sup>th</sup> November sowing date found comparatively better in terms of increasing plant height (8.51%), tiller (14.90%) and dry matter accumulation (13.38%), compared to 30<sup>th</sup> December sowing. The maximum

growth rate of the plants under the 30<sup>th</sup> November sowing might be due to plants get more time for their growth and development compared to late sowing which has short growth and a short time period.

Among varieties, HD 2967 exhibited Maximum plant height (110.40 cm), number of tillers (520.08 m<sup>-2</sup>), dry weight (1070.85 g m<sup>-2</sup>) compared to K-307. The variety HD 2967 recorded an increase in terms of plant height (7.06%), number of tillers (2.8%), dry matter accumulation (25.71%) compared to the variety K-307.

Probably because of the better genomics of the variety HD 2967, which excels in terms of growth and development, especially in particular climatic conditions compared to other varieties.

These results are in conformity with the findings of Anil Kumar Netam *et al.* (2020) <sup>[10]</sup>, Bidusi Tripathy *et al.* (2020), Sandeep Kumar Singh *et al.* (2020), Mohammad Yusuf *et al.* (2019).

#### Effect of treatments on yield attributing characters of wheat varieties

The Data pertaining to yield attributing character revealed that 30<sup>th</sup> November sowing was found significantly better than 15<sup>th</sup> December and 30<sup>th</sup> December. The increment recorded under different yield attributes varied to the tune in length of ear (5.72 %), ear weight (19.20%), grains per ear-1 (15.75%), test weight (7.50%), under 30<sup>th</sup> November sowing compared to 30<sup>th</sup> December sowing. The better yield attributes under 30<sup>th</sup> November sowing is probably due to the good vegetation at early sowing compared to 30<sup>th</sup> December sowing.

The variety HD 2967 recorded better yield attributes varied the tune in the length of the ear length (9.40%), ear weight (10.75%), grain ear-1 (9.22%), and test weight (2.77%) compared to the variety K-307. Similar findings were also reported by Charanjeet Kaur (2017), Bashir *et al.* (2016), Kaur *et al.* (2015), Dwivedi *et al.* (2015) and Shirpurkar *et al.* (2007).

#### Effect of treatments on yield of wheat varieties

The data pertaining to the yield of wheat was significantly influenced by different dates of sowing and varieties. The biological yield of wheat increased 5.38% and 13.93 % under 30<sup>th</sup> November compared to 15<sup>th</sup> December and 30<sup>th</sup> December sowing.

Similarly, grain yield increased 44.68 % in the 30<sup>th</sup> November sowing compared to 30<sup>th</sup> December sowing respectively. The Straw yield increased 17.73% under the 30<sup>th</sup> November sowing compared to the 30<sup>th</sup> December sowing. The harvest index was also increased 13.98 % under 15<sup>th</sup> November compared to 30<sup>th</sup> December sowing.

Under different varieties HD 2967 produced maximum biological yield (11.32%), grain yield (27.52%), straw yield (21.22%), and harvest index (28.94%) compared to K-307. It might be due to better genomics characters as well as better growth characteristics and yield attributes of the variety HD 2967 achieved under favourable climatic conditions.

Similar findings were reported by Amarjeet *et al.* (2020), Mohammad Yusuf *et al.* (2019), Chetan Lal (2019), Kumar *et al.* (2015).

**Table 1:** Effect of weather parameters, date of sowing on performance of wheat varieties

Treatment	Growth Attributes				Yield Attributes				Yield			
	Plant population 2 (m)	Plant height (cm)	Til2lers (m)	Dry matter accumulation (g/m <sup>2</sup> )	Length of ear (cm)	Ear weight (g)	Grains ear-1	Test weight (g)	Biological yield (q/ha-1)	Grain yield (q/ha-1)	Straw yield (q/ha-1)	Harvest Index (%)
<b>Date of sowing</b>												
15th Nov	123.93	111.925	537.875	1152.20	9.60	2.11	53.55	39.31	115.16	44.94	70.22	39.05
30th Nov	122.65	108.919	519.500	1060.03	9.40	1.98	48.70	37.31	105.90	37.20	68.70	35.15
15th Dec	122.44	103.056	468.125	920.169	9.08	1.77	46.26	36.56	90.71	31.06	59.64	34.26
SE(d)±	0.490	0.638	1.275	10.169	0.049	0.016	0.479	0.156	1.001	0.663	0.420	0.320
CD (P=0.05)	NS	1.592	3.180	25.367	0.122	0.041	1.196	0.390	2.496	0.500	1.047	0.799
<b>Varieties</b>												
HD 2733	124.058	107.717	512.417	1048.04	9.17	1.98	49.72	37.85	103.10	40.15	62.94	38.67
K-307	123.091	103.175	505.750	973.992	8.91	1.92	47.37	37.60	97.30	35.31	61.99	36.09
K-9107	122.681	110.733	495.750	1084.05	9.75	1.86	49.18	37.44	108.32	33.17	75.15	30.51
HD 2967	122.208	110.242	520.083	1070.45	9.61	2.06	51.74	38.46	106.98	42.30	64.68	39.34
SE(d)±	0.376	0.520	2.078	8.897	0.058	0.018	0.361	0.129	0.944	0.707	0.420	0.413
CD (P=0.05)	NS	1.073	4.286	18.353	0.120	0.036	0.744	0.267	1.947	1.459	1.047	0.852
<b>Interaction</b>												
<b>V X D</b>												
SE(d)±	0.651	0.901	3.599	15.410	0.101	0.030	0.625	0.224	1.635	1.225	0.805	0.716
C.D. (P=0.05)	NS	NS	NS	NS	NS	NS	1.408	NS	NS	2.676	1.753	1.541
<b>D X V</b>												
SE(d)±	0.747	1.008	3.368	16.778	0.100	0.031	0.723	0.249	1.734	1.251	0.814	0.698
C.D. (P=0.05)	NS	NS	NS	NS	NS	NS	1.628	NS	NS	2.733	1.772	1.503

## References

1. Abad AL, Loveras J, Michlena A. Nitrogen fertilization and foliar urea effects on Durum wheat yield and quality and on residual soil nitrate in irrigated Mediterranean conditions. *Field Crops Research*. 2004;87:257-269.
2. Afzal M, Nazir MS. Response of two semi dwarf wheat varieties to sowing dates. *Journal of Agricultural Research*. 1986;24:110-114.
3. Agrawal DK, Nath S. Effect of climatic factor and date of sowing on wheat crop in Allahabad condition, Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(9):1776-1782.
4. Agrawal KK, Sanker U, Upadhyay AP, Gupta VK. Accumulated heat unit requirements for different phenophases of wheat (*Triticum aestivum*) cultivars as influenced by sowing dates at Jabalpur. *Journal of Agro meteorology*. 1999;1(2):173-176.
5. Agrawal PK, Sinha SK. Effect of probable increase in carbon dioxide and temperature on wheat yield in India. *Journal of Agricultural Meteorology*. 1993;48(5):811-814.
6. Ahmad Z. Effect of irrigation and cultivar on the growth and yield of wheat. M.Sc. (Hons.) Agri. Thesis, Univ. Agri., Faisalabad, Pakistan, 1991.
7. Ahuja KN, Lal RB, Ashok Kumar, Kumar A. Effect of seed rate, date and method of sowing on growth and yield of wheat. *Annals of Agricultural Research*. 1996;17:190-192.
8. Alexander B, Hayman P, Mc Donald G, Talukder HASM Gill G. () Characterizing the risk of heat stress on wheat in South Australia: Meteorology, climatology and the design of a field heating chamber. In: Dove, H. (Ed.), *Food Security from Sustainable Agriculture: Proceedings of 15th Agronomy Conference*, Lincoln, New Zealand; c2010. p. 15-19.
9. Ali MH. Irrigation-yield response factor of winter wheat for different growth phases. *Journal of Agrometeorology*. 2009;11(1):9-14.
10. Anil Kumar Netam, Upendra Kumar Nag, Chainu Ram Netam. Growth and Yield of Wheat (*Triticum aestivum* L.) Varieties as Influenced by Different Sowing Dates under Bastar Plateau Zone of Chhattisgarh. *Int. J Curr. Microbiol. App. Sci*. 2020;9(6):2161-2169.