



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2021; 4(2): 186-189

Received: 23-09-2021

Accepted: 22-11-2021

**Jadhav KK**

M.Sc. Agri. Scholar, Agronomy,  
College of Agriculture, Pune,  
Maharashtra, India

**Kashid NV**

Associate Professor, Agronomy,  
College of Agriculture, Pune,  
Maharashtra, India

**Avte SB**

M.Sc. Agri. Scholar, Soil science  
and agricultural chemistry, College  
of Agriculture, Parbhani,  
Maharashtra, India

**Halge SS**

M.Sc. Agri. Scholar, Agronomy,  
College of Agriculture, Pune,  
Maharashtra, India

## Effects of planting methods on yield attributes and yield of paddy (*Oryza sativa* L.)

**Jadhav KK, Kashid NV, Avte SB and Halge SS**

DOI: <https://doi.org/10.33545/2618060X.2021.v4.i2a.158>

### Abstract

An experiment entitled, “Effects of planting methods on growth and yield of paddy (*Oryza sativa* L.)” was carried out during *Kharif*, 2020 at Agricultural Research Station Farm, VadgaonMaval, Tal. Maval, Dist. Pune to investigate effects of planting methods on yield attributes and yield of paddy (*Oryza sativa* L.). The field experiment was laid out in Randomized Block Design (RBD) with three replications. There were eight treatments comprising of different sowing methods of rice *viz.*, T<sub>1</sub>-Sowing as direct seeded rice (DSR) at 22.5cm by bullock drawn seed drill, T<sub>2</sub>-Sowing as direct seeded rice (DSR) at 30cm by bullock drawn seed drill, T<sub>3</sub>-Sowing by dibbling method at 20 x 15 cm<sup>2</sup>, T<sub>4</sub>-Direct sowing of rice by tractor operated mechanical seed drill, T<sub>5</sub>-Direct sowing of rice by ‘Saguna Rice Technique’(SRT), T<sub>6</sub>-Direct sowing of rice by ‘Modified Direct Seeded Rice Technique’(MDSRT), T<sub>7</sub>-University recommended method (Four Point Agro-Technology or Char Sutri Method) and T<sub>8</sub>-Farmer’s practice-Conventional transplanting method. The periodical observations on yield contributing characters and yield were recorded to assess the treatment effects. Results revealed that planting methods had significant effects on yield attributes and yield of paddy. The yield attributing characters *viz.*, length of spike (cm), number of spikes per plant, number of grains per spike, grain weight per spike (g) and thousand grain weight (g) were found higher with T<sub>6</sub>-Direct sowing of rice by ‘Modified Direct Seeded Rice Technique’ (MDSRT) than other planting methods. The corresponding values were 26.3 cm, 8.3, 161.4, 3.12 g and 23.0 g respectively. The grain yield (63.74 q ha<sup>-1</sup>) and straw yield (68.88 q ha<sup>-1</sup>) were significantly superior in the treatment T<sub>6</sub>-Direct sowing of rice by ‘Modified Direct Seeded Rice Technique’ (MDSRT) (63.74 q ha<sup>-1</sup>) than rest of the sowing methods except it was at par with the treatment T<sub>7</sub>-University recommended method (Four Point Agro-technology) having grain yield (55.60 q ha<sup>-1</sup>) and straw yield of paddy (60.10 q ha<sup>-1</sup>) respectively.

**Keywords:** Planting methods, DSR, MDSRT, paddy, yield attributes and yield

### Introduction

Rice (*Oryza sativa* L.) is one of the most ancient crops being cultivated in 117 countries, hence called as “Global grain”. It is the staple cereal food grain of majority of India’s over one billion population, contributes to nearly 44 per cent of total food grain production. Rice feeds more people over a longer period of time than any other crop. Rice has been documented in the history books as a source of food and for tradition as well since 2500 B.C. Beginning in China and the surrounding areas, its cultivation spread throughout Sri Lanka and India. Global demand for food is rising because of population growth, increasing affluence and changing dietary habits. The UN/FAO forecasts that global food production will need to increase by over 40 per cent by 2030 and 70 per cent by 2050. Yet globally, water is anticipated to become scarce and there is increasing competition for land, putting added pressure on agricultural production. In addition, climate change will reduce the reliability of food supply through altered weather patterns and increased pressure from pests and diseases. Rice along with wheat form the bedrock of Indian food security and to meet the country’s stated goal of ensuring food for all, farmer will have to produce more rice from lesser land, using less water, energy and other inputs and keeping in harmony with the fragile environment.

The production of conventional puddle transplanted rice faces severe constraints because of water and labour scarcity and climatic changes (Pathak *et al.*, 2011)<sup>[9]</sup>.

**Corresponding Author:**

**Jadhav KK**

M.Sc. Agri. Scholar, Agronomy,  
College of Agriculture, Pune,  
Maharashtra, India

Imminent water crisis, water-demanding nature of traditionally cultivated rice and climbing labour costs ramble the search for alternative management methods to increase water productivity, system sustainability and profitability. Direct seeded rice (DSR) technique is becoming popular nowadays because of its low-input demanding nature. It offers a very exciting opportunity to improve water and environmental sustainability. It is a feasible alternative to conventional puddled transplanted rice with good potential for saving water, mitigating greenhouse gas emissions and adapting to climatic risks and the yield can be comparable with that of transplanted rice if the crop is properly managed (Kumar and Ladha, 2011) [6]. It involves sowing pre-germinated seeds into a puddle soil surface (wet seeding), standing water (water seeding) or dry seeding into a prepared seedbed (dry seeding). Recently there is trend towards direct seeded rice because of labour and water scarcity. Although the development of suitable varieties and agronomic packages for promoting direct-seeded rice is under way (Pathak *et al.*, 2011) [9], so far no variety has been developed that possess traits specifically needed to high yield under dry direct-seeded conditions, particularly for rainfed systems that may be prone to drought and low fertility.

In the North-Western Indo-Gangetic Plains (IGP), transplanted rice is predominantly cultivated. Transplanting requires at least 25 ha-cm of water for puddling operation, which creates a dense clay layer in the sub-soil to prevent seepage losses. The crop requires about  $130 \pm 10$  ha-cm of irrigation in addition to the adoption of suitable variety and application of a recommended dose of fertilizers to realize yield levels of about  $6 \pm 2$  t/ha. Generally, about 40% of all irrigation water goes to paddy cultivation in the region. It is estimated that flooded rice fields produce about 10% of global methane emissions. Also, injudicious use of nitrogenous fertilizers is a common feature in paddy cultivation which is a source of nitrous oxide emissions.

The current practice of excessive exploitation of groundwater has led to a decline in the quality of natural resources i.e., land and water. Researchers have developed suitable drilled paddy alternatives to transplanted paddy. In drilled paddy cultivation, raising of nursery for transplantation is done away with. The farmer can avoid the major problem faced i.e., labour shortage for transplanting due to peak demand. In case of delay in monsoon or shortage of water, a drilled paddy gives the farmer flexibility to take up direct sowing of paddy with a suitable duration variety to fit into the left over a season. Drilled paddy consumes relatively less water compared to transplanted flooded rice. Energy demand for pumping of irrigation water is also less and saving can be much higher during deficit rainfall situations compared to transplanted rice. Direct sowing can be practiced for cultivating both coarse rice and basmati rice wherever feasible. Drilled paddy with reduced tillage is an efficient resource conservation technology that holds great promise in the Indo-Gangetic Plain because of the following advantages.

- Saving in water up to 25% in DSR
- Saving in energy up to 27% of diesel as pumping energy is saved for field preparation, nursery raising, puddling and reduced frequency of applying irrigation water
- Saving of 35-40 man days / ha
- Enhanced fertilizer use efficiency due to the placement of fertilizer in the root zone
- Early maturity of crops by 7-10 days helps in the timely sowing of succeeding crops (De Datta, 1986) [1].
- Reduction in methane emissions and global warming potential.
- Little disturbance to soil structure

- Enhanced system productivity

Agronomic manipulations *viz.*, planting geometry, density may be advantageous for achieving the potential yield of paddy. The optimum planting geometry differed for different planting methods. Hence, determination of suitable establishment method for harnessing the potential yield of different paddy varieties needs the critical investigations. On the other hand, non-availability of irrigation water, insufficient labour and high wages during the peak period of farm operation invariably delay planting of paddy. To mitigate this problem, many rice farmers are switching to direct seeding of paddy. Direct seeding can reduce the labour requirement, may reduce methane gas emission, shorten the duration of crop by 7 to 10 days and provide comparable grain yield to transplanting (De Datta, 1986) [1].

## Materials and Methods

The field experiment was conducted during *Kharif*, 2020 at Agricultural Research Station Farm, Vadgaon Maval, Tal. Maval, Dist. Pune. to study the effects of planting methods on yield attributes and yield of paddy (*Oryza sativa* L.)". The soil of experimental field was clay loam in texture, moderately alkaline in reaction (pH 7.74) with low available nitrogen ( $239 \text{ kg ha}^{-1}$ ), medium available phosphorus ( $16 \text{ kg ha}^{-1}$ ) and high available potassium ( $389 \text{ kg ha}^{-1}$ ).

The experiment was laid out in Randomized Block Design (RBD) with eight treatments and three replications. The paddy variety VDN-99-29 (*Phule Samruddhi*) is cultivated using eight methods *viz.*, T<sub>1</sub>-Sowing as direct seeded rice (DSR) at 22.5cm by bullock drawn seed drill, T<sub>2</sub>-Sowing as direct seeded rice (DSR) at 30cm by bullock drawn seed drill, T<sub>3</sub>-Sowing by dibbling method at  $20 \times 15 \text{ cm}^2$ , T<sub>4</sub>- Direct sowing of rice by tractor operated mechanical seed drill, T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique'(SRT), T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique'(MDSRT), T<sub>7</sub>-University recommended method (Four Point Agro-Technology or Char Sutri Method) and T<sub>8</sub>-Farmer's practice-Conventional transplanting method. The gross plot size was 3.60 m x 3.00 m and net plot size was different as per treatments.

RDF of 100:50:50 NPK  $\text{kg ha}^{-1}$  was applied during conduct of experiment. Out of which half dose of N and full dose of P and K are applied as basal dose at the time of sowing i.e., 50:50:50 NPK  $\text{kg ha}^{-1}$ . While remaining dose of N is split into two and applied after each hand weeding. For the treatments T<sub>6</sub>-Direct sowing of rice 'Modified Direct Seeded Rice Technique' (MDSRT) and T<sub>7</sub>-University recommended method (Four Point Agro-Technology), the N and P fertilizers are given through 170 kg Urea-DAP briquettes (60:30:00) and 50 kg K<sub>2</sub>O as straight fertilizer per hectare. Gap filling was done at 15<sup>th</sup> days after sowing, two manual weedings were done throughout the rice growing period.

For recording growth observations, five plants were selected randomly from each net plot. The selected plants were labeled and were marked by fixing pegs near them. All the observations on growth and yield were recorded on these plants. The crop from each net plot was harvested separately at maturity, labeled and tied in bundles according to treatments. The produce of each plot was threshed separately and weight of grain and straw taken separately. The experimental data was statistically analyzed by using analysis of variance in the randomized block design (RBD) (Panse and Sukhatme, 1985) [8].

## Result and Discussion

### Yield attributes

The yield attributes of paddy viz., length of spike plant<sup>-1</sup>, number of spikes plant<sup>-1</sup>, number of grains per spike, grain weight per spike (g) and test weight (g) were significantly influenced by different planting methods (Table 1). The treatment T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded significantly maximum mean length of spike (26.3 cm) which was at par with the treatment T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique' (SRT) having mean length of spike (25.7 cm), T<sub>7</sub>-University recommended method (Four Point Agro-technology) having mean length of spike (24.2 cm), T<sub>3</sub>-Sowing by dibbling method at 20x15 cm<sup>2</sup> having mean length of spike (22.8 cm) and T<sub>4</sub>-Direct sowing of rice by tractor operated mechanical seed drill (22.2 cm). Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded the maximum length of spike throughout the growth period, this might be due to favourable climatic condition for better seedling establishment and further growth and development of *kharif* paddy crop. Similar results were reported by Kaur and Mahal (2016) as low spacing severely affected the yield attributing characters. The treatment T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded significantly maximum mean number of spikes plant<sup>-1</sup> (8.3) than rest of all the treatments except treatments T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique' (SRT) having number of spikes plant<sup>-1</sup> (8.2) and T<sub>7</sub>-University recommended method (Four Point Agro-technology) having number of spikes plant<sup>-1</sup> (7.8), where those treatments found statistically at par with the best one. Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded the maximum numbers of spikes plant<sup>-1</sup> throughout the growth period, this might be due to better seedling establishment and favourable climatic condition for growth and development of *kharif* paddy crop. Similar results were reported by Kaur and Singh (2016). The treatment T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded significantly maximum number of grains spike<sup>-1</sup> (161.4) than rest of all the

treatments except it was at par with the treatment T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique' (SRT) having number of grains spike<sup>-1</sup> (155.3) and T<sub>7</sub>-University recommended method (Four Point Agro-technology) having number of grains spike<sup>-1</sup> (137.6). The treatment T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded significantly maximum grain weight per spike (3.12 g) than rest of all the treatments except it was at par with the treatment T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique' (SRT) having grain weight per spike (3.11 g), T<sub>7</sub>-University recommended method (Four Point Agro-technology) having grain weight per spike (2.85 g), T<sub>3</sub>-Sowing by dibbling method at 20 x 15 cm<sup>2</sup> having grain weight per spike (2.68 g) and T<sub>4</sub>-Direct sowing of rice by tractor operated mechanical seed drill having grain weight per spike (2.61 g). Similar results were reported by Thakur *et. al.* (2010). The treatment T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) recorded significantly maximum test weight (23 g) which was at par with the treatment T<sub>5</sub>-Direct sowing of rice by 'Saguna Rice Technique' (SRT) having test weight (22.5 g), T<sub>7</sub>-University recommended method (Four Point Agro-technology) having test weight (22.8 g), T<sub>3</sub>-Sowing by dibbling method at 20 x 15 cm<sup>2</sup> having test weight (22.3 g) and T<sub>4</sub>-Direct sowing of rice by tractor operated mechanical seed drill having test weight (22.0 g). The treatment of direct sowing of rice 'Modified Direct Seeded Rice Technique' (MDSRT) recorded maximum test weight (23.0) which was significantly superior over rest of all the cultivation methods. Similar results were also reported by Hundal and Kaur (1995). The yield attributing characters viz., length of spike (cm), number of spikes per plant, number of grains per spike, grain weight per spike (g) and thousand grain weight (g) were found higher with direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) than other planting methods. The lowest length of spike (cm), number of spikes per plant, number of grains per spike, grain weight per spike (g) and thousand grain weight (g) were recorded in the treatment T<sub>8</sub>-Farmer's practice- Conventional transplanting method.

**Table 1:** Length of spike (cm), Number of spikes per plant, Number of grains per spike, Grain weight per spike (g) and Thousand grain weight (g) of paddy as affected by different treatments

Tr. No.	Treatments	Length of spike (cm)	Number of spikes per plant	Number of grains per spike	Grain weight per spike (g)	Thousand grain weight (g)
T <sub>1</sub>	Sowing as direct seeded rice (DSR) at 22.5 cm by bullock drawn seed drill.	21.0	4.4	118.4	2.47	21.2
T <sub>2</sub>	Sowing as direct seeded rice (DSR) at 30 cm by Bullock drawn seed drill	21.8	5.8	122.7	2.56	21.7
T <sub>3</sub>	Sowing by dibbling method At 20x15 cm <sup>2</sup>	22.8	5.8	128.3	2.68	22.3
T <sub>4</sub>	Direct sowing of rice by Tractor operated mechanical seed drill.	22.2	5.9	131.0	2.61	22.0
T <sub>5</sub>	Direct sowing of rice by 'Saguna Rice Technique' (SRT)	25.7	8.2	155.3	3.11	22.5
T <sub>6</sub>	Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT)	26.3	8.3	161.4	3.12	23.0
T <sub>7</sub>	University recommended method (Four Point Agro-technology)	24.2	7.8	137.6	2.85	22.8
T <sub>8</sub>	Farmer's practice- Conventional transplanting method	13.0	3.3	66.2	1.32	18.25
	S.E.m.±	1.36	0.43	8.67	0.17	1.50
	C.D.at5%	4.14	1.32	26.31	0.52	4.57
	General Mean	22.10	6.21	127.60	2.59	21.70

### Yield studies

The grain yield and straw yield (q ha<sup>-1</sup>) of paddy were significantly influenced by different treatments (Table 2). The

grain yield of paddy was influenced significantly due to different cultivation methods. The grain yield of paddy (q ha<sup>-1</sup>) was found superior in the treatment T<sub>6</sub>-Direct sowing of rice by 'Modified



Direct Seeded Rice Technique' (MDSRT) ( $63.74 \text{ q ha}^{-1}$ ) than rest of the sowing methods which was at par with the treatment T<sub>7</sub>-University recommended method (Four Point Agro-technology) having grain yield ( $55.60 \text{ g}$ ). The lowest grain yield ( $25.83 \text{ q ha}^{-1}$ ) was produced by the treatment T<sub>8</sub>-Farmer's practice-Conventional transplanting method. The treatment of direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) (T<sub>6</sub>) recorded maximum grain yield ( $63.74 \text{ q ha}^{-1}$ ) which was significantly superior over rest of all the cultivation methods due to proper spacing, better establishment, favourable microclimatic conditions and less attack of diseases and pest. Similar results were reported by Gunri *et al.* (2004) <sup>[11]</sup> and Christian (2017). The straw yield of paddy was influenced significantly due to different cultivation

methods. The straw yield was found significantly superior ( $68.88 \text{ q ha}^{-1}$ ) under the treatment of T<sub>6</sub>-Direct sowing of rice by 'Modified Direct Seeded Rice Technique' (MDSRT) than rest of all other planting methods which was at par with the treatment T<sub>7</sub>-University recommended method (Four Point Agro-technology) having straw yield ( $60.10 \text{ q ha}^{-1}$ ). The lowest straw yield was obtained in the treatment T<sub>8</sub>-Farmer's practice-Conventional transplanting method ( $28.17 \text{ q ha}^{-1}$ ). The treatment of direct sowing of rice Modified Direct Seeded Rice Technique (MDSRT) recorded maximum straw yield ( $68.88 \text{ q ha}^{-1}$ ) which was significantly superior over rest of all the cultivation methods due to proper spacing, better establishment, favourable microclimatic conditions and less attack of diseases and pest. Similar result was found by Muhammad (2014) <sup>[7]</sup>.

**Table 2:** Grain yield ( $\text{q ha}^{-1}$ ) and straw yield ( $\text{q ha}^{-1}$ ) of paddy as affected by different treatments

Tr. No.	Treatments	Grain yield ( $\text{q ha}^{-1}$ )	Straw yield ( $\text{q ha}^{-1}$ )
T <sub>1</sub>	Sowing as direct seeded rice(DSR) at 22.5cm By bullock drawn seed drill.	48.78	53.74
T <sub>2</sub>	Sowing as direct seeded rice (DSR) at 30 cm by Bullock drawn seed drill	49.86	55.24
T <sub>3</sub>	Sowing by dibbling method at $20 \times 15 \text{ cm}^2$	52.78	58.13
T <sub>4</sub>	Direct sowing of rice by tractor operated Mechanical seed drill.	51.42	56.68
T <sub>5</sub>	Direct sowing of rice by 'Saguna Rice Technique'(SRT)	48.66	54.07
T <sub>6</sub>	Direct sowing of rice by 'Modified Direct Seeded Rice Technique'(MDSRT)	63.74	68.88
T <sub>7</sub>	University recommended method (Four Point Agro-technology)	55.60	60.10
T <sub>8</sub>	Farmer's practice-Conventional transplanting Method	25.83	28.17
	S.E.m. <sub>±</sub>	2.91	3.18
	C.D.at5%	8.84	9.67
	General Mean	49.58	54.37

## Conclusion

Based on the present investigation, it is concluded that the direct sowing of rice with 'Modified Direct Seeded Rice Technique' (MDSRT) and transplanting of rice with university recommended method (Four Point Agro-technology) recorded higher yield attributes and yield of paddy than rest of all the treatments. Thus, from economic and labour constraint point of view, direct sowing of rice with 'Modified Direct Seeded Rice Technique' (MDSRT) or transplanting of rice with university recommended method (Four Point Agro- technology) are advisable.

## Acknowledgement

The authors are thankful to Agronomy Section, College of Agriculture, Pune - 411005, Maharashtra, India for providing necessary facilities to undertake the field experiment.

## References

- DeDatta SK. Technology development and the spread of direct seeded flooded rice in South East Asia. Fertilizer Research Journal. 1986;9:171-186.
- Sumekar Y, Widayat D, Riswandi D, Kurniadie D. Effects of metsulfuron methyl herbicide doses on weed suppression, growth and yield of rice (*Oryza sativa* L.). Int. J Agric. Food Sci. 2020;2(1):18-23. DOI: 10.33545/2664844X.2020.v2.i1a.30
- Hundal SS, Kaur P. Effect of environmental stresses on potential production of major cereal crops in Punjab In. Proc. Int. Conf. Sust. Agri. Env. held at Haryana Agri. Univ. Hisar. 1995;9(2):11-12.
- Kaur J, Mahal SS. Irrigation scheduling in direct seeded rice (*Oryza sativa* L.). An Agricultural Reviews. Indian Journal of Agronomy. 2015;36(3):208-217.
- Kaur S, Singh S. Influence of plant densities on growth and yield of direct Seeded rice (*Oryza sativa* L.). Agricultural Research Journal. 2016;53(3):594-596.
- Kumar V, Ladha JK. Direct seeding of rice: recent developments and future research needs. Advances in Agronomy. 2011;111:297-413.
- Muhammad AI. Productivity and Quality of Direct Seeded Rice under Different Types of Mulches and Planting Patterns: A Review. American-Eurasian Journal of Agricultural and Environmental Science. 2014;14(11):1240-1247.
- Panse VG, Sukhatme PV. Statistical methods for Agriculture work. 4th Edn. ICAR, New Dehli. 1985;145-156.
- Pathak H, Tewari AN, Sankhyani S, Dubey DS, Mina U, Singh VK, *et al.* Direct-seeded rice: Potential, performance and problems - A review. Current Advances in Agricultural Sciences. 2011;3(2):77-88.
- Thakur AK, Uphoff N, Antonyu E. An assessment of physiological effects of system of rice intensification (SRI) practices compared with recommended rice cultivation practices in India. Experimental Agriculture. 2010;46(1):77-98.
- Gunri SK, Pal SK, Choudhary A. Effect of nitrogen application and spacing on yield of rice in foot-hill soil of West Bengal. Indian Journal of Agronomy. 2004;49(4):248-250.