



International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

www.agronomyjournals.com

2022; 5(2): 199-206

Received: 02-11-2022

Accepted: 07-12-2022

Hayilu Gichile

Haramaya University, P. O. Box
138, Haramaya, Ethiopia

Review on methods of maize breeding for tolerance or resistance of *Striga hermonthica*

Hayilu Gichile

DOI: <https://doi.org/10.33545/2618060X.2022.v5.i2c.176>

Abstract

Striga hermonthica is a major limiting factor in maize cultivation in sub-Saharan Africa. Based on environmental factors such as soil fertility, climatic conditions and genotype, this can result in a yield loss of up to 100%. The aim of this article is to review maize breeding methods for resistance/tolerance to *Striga hermonthica* and to highlight the efforts made to overcome yield reduction due to *Striga hermonthica*. The International Institute of Tropical Agriculture (IITA) has made significant progress in developing several early inbred maize lines, open-pollinated varieties and hybrids containing *Striga* resistance genes. However, the levels of *striga* resistance of the available precocious inbreds and hybrids are not as high as desired. This is because the *Striga* seed bank is added to the soil after each growing season. There is therefore a need for increased *Striga* resistance in the available genotypes. Since resistance to *S. hermonthica* in maize is regulated by many genes, breeding for *Striga*-resistant cultivars using traditional approaches has been less effective and time-consuming. The identification of QTLs associated with *Striga* resistance/tolerance would facilitate the rapid development of *Striga* resistant/tolerant maize genotypes.

Keywords: Maize, parasite, *striga hermonthica*, *striga* resistance, *striga* tolerance

1. Introduction

Maize (*Zea mays* L.) is the world's most widely grown crop and the most important staple food in Sub-Saharan Africa (SSA) (FAO, 2017) ^[30]. It is the most important food crop on the African continent in terms of acreage and total grain production (Reynolds *et al.*, 2015) ^[66]. Corn is the main food source for over 300 million people. It is a major source of carbohydrates, proteins, lipids and vitamins for millions of people in the region (Badu-Apraku *et al.* 2017) ^[19]. A high annual consumption of 79 kg per capita in Africa (De Groote, 2002). Corn can be exploited in a variety of ways, with all parts of the plant such as kernel, cob, stalk, leaves and tassel having economic value. The grain is used as food or fermented to make a variety of beverages and beers (Anami *et al.*, 2009) ^[10].

However, corn production is still below the crop's potential yields due to biotic and abiotic constraints, leading to food insecurity (FAO, 2017) ^[30]. Among parasitic weeds, the root hemiparasite *S. hermonthica* is the most devastating and dominant constraint on maize production (Khan *et al.*, 2014) ^[44]. The extent of damage depends on the timing and extent of infection. Can cause up to 100% in high infestation (Amusan *et al.* 2008) ^[89]. Losses due to *Striga* are estimated at \$7 billion annually. Today, the weed affects over 100 million farmers (Spallek *et al.*, 2013) ^[76]. *Striga hermonthica* (giant witch's herb), a parasitic plant native to Ethiopia and Sudan (Amy. B *et al.*, 2011) ^[9]. A separate survey conducted in 1997 found that *Striga hermonthica* is the most widespread parasitic weed species in Ethiopia and the overall incidence rate of *Striga* of the 310 maize fields surveyed was 41%.

Using cultivars that are tolerant and resistant to *Striga* species has been recommended as the most practical approach for resource-poor smallholders (Menkir, 2006) ^[52]. Since 1982, scientists at IITA have been breeding maize for tolerance and resistance to *S. hermonthica*. Genetic resistance in the host plant is central to the success of integrated control measures to minimize the *Striga* threat. Decades of *Striga* research efforts at the International Institute of Tropical Agriculture (IITA) Ibadan have recently resulted in the successful release and registration of *Striga*-resistant maize inbred lines from diverse genetic backgrounds.

Corresponding Author:

Hayilu Gichile

Haramaya University, P. O. Box
138, Haramaya, Ethiopia

Planting *Striga*-resistant maize varieties is currently considered the best control strategy and is easy to apply or deploy, especially in combination with other management practices (Gasura, E., *et al.*, 2019) [31]. Resistance to *S. hermonthica* parasitism is mainly attributed to the low production of *Striga* germination stimulants by the host plant (Rodenburg, J. *et al.*, 2006) [68]. When breeding for *S. hermonthica* resistance in maize, a combination of these resistance mechanisms is desirable to achieve effective and durable resistance (Gasura, E., *et al.*, 2019) [31]. The slow evolution and spread rate of *Striga*-resistant genotypes is largely due to the complex genetics of resistance as well as limited knowledge of the specific mechanisms associated with resistance to *Striga* (Amusan, *et al.*, 2008) [89].

Resistance to *S. hermonthica* in maize is polygenic and is significantly influenced by the environment (Redinbaugh M, Pratt, R., 2009) [65]. Because of this, breeding *Striga*-resistant cultivars using conventional approaches has been less effective and time-consuming (Ejeta and Gressel 2007) [26]. Marker assisted breeding utilizes genotypic data in identifying genotypes possessing desirable alleles using linked genetic markers. QTL mapping approaches are important genomic tools used in analyzing the genetic architecture of complex traits (Awata, LAO *et al.*, 2020; Redinbaugh, M.; Pratt, R., 2009) [13, 65] as well as in identifying genetic linkage through broad Genotyping a panel of germplasm with contrasting phenotypes in different environments (Srivastava, R.K.; *et al.* 2020) [77]. Genetic engineering is another option for breeding *Striga*-resistant strains. It offers an effective alternative to quickly establish resistance to *Striga* spp. to achieve, based on reports of other parasitic plants.

Although many efforts have been made by plant breeders (IITA) resulting in the development and release of several *Striga*-resistant/tolerant maize inbred lines, limited success has been achieved in controlling *Striga* in smallholder fields in Africa. This is mainly because the *Striga hermonthica* has a highly specialized life cycle synchronized with host growth, a breeding behavior that maintains tremendous genetic variability, the ability to parasitize a wide range of hosts, seed longevity in soil, and the ability to inflict the most damage to the host before emerging above ground (Ejeta, 2007) [26]. Therefore, it is very important to develop strategies integrating different breeding techniques for the development of a *S. hermonthica* resistant maize variety that would help reduce losses due to this parasite infestation suffered by maize farmers in sub-Saharan Africa, including Ethiopia. Therefore, this review examines the different approaches of maize breeding for resistance (tolerance) to parasitic weeds called *Striga hermonthica* and the aim of this paper was to review on methods of maize breeding for resistance/tolerance to *Striga hermonthica* and highlight efforts made to overcome yield reduction due to *Striga hermonthica*.

2. Literature Review

2.1 Origin, occurrence and distribution of *Striga hermonthica*

Studies have shown, among genus *Striga*, 80% are found in Africa while the rest occur in Asia and the United States. Many reports indicated that, *S. hermonthica* originated from the Nubian hills of Sudan and Semien mountains of Ethiopia (Atera *et al.*, 2014) [12]. Currently most parts of Africa are infested with *S. hermonthica* parasite with heavy infestation in some parts of east and West Africa. Moderate to light infestation is shown in central Africa and Southern Africa (fig).




Fig 1: Show the striga infestation

S. hermonthica can tolerate very different environments and adapt very quickly to different hosts. The study showed that geographic distance played a greater role in the genetic differentiation of *S. hermonthica* populations than host specialization. It has been shown that genetic differentiation based on host specificity is greater between the millet and sorghum strains than between any maize strains. In Ethiopia, the Rift Valley is a potential geographic barrier to the spread of *S. hermonthica* (Amy. B. *et al.* 2011) [9]. A second source of *Striga* seed dispersal is contamination of grain kernels with parasitic plant seeds. This is because crops in contaminated fields are threshed onto the ground using oxen and donkeys. The oxen and donkeys provide a secondary source of seed dispersal.

2.2 *Striga* lifecycle and its economic importance

The lifecycle of *Striga* is synchronized to that of its host and involves mechanisms that coordinate the lifecycle of the parasite and that of the host (Bouwmeester *et al.*, 2003) [22]. The *Striga* life cycle generally involves: germination, host attachment, formation of haustoria, penetration, and establishment of vascular connections, nutrients accumulation, flowering and production of seeds. *Striga* seeds only germinate in presence of certain hormones known as strigolactones, produced by the host and in other cases non-host species (Keyes *et al.*, 2007) [90].




Fig 2: Major stages in the life cycle of *Striga* of development (Bouwmeester *et al.*, 2003) [22]

The economic losses caused by *Striga* spp. are enormous. This is a parasite found mainly on tropical grain crops such as corn, sorghum, pearl millet and upland rice (Press *et al.*, 2001). It causes reduced growth for the host because it disrupts its

photosynthesis and uses its nutrients, causing a deficit (Joel, 2007) [40]. The weeds consume a larger portion of the host plant's solutes, resulting in wilting and early death of the plant (Ruyter-Spira *et al.*, 2011) [74]. It is estimated that in sub-Saharan Africa alone there is an annual yield loss of more than US\$7 billion due

to infestation with *Striga* spp. (Spallek *et al.*, 2013) [76]. The amount of yield loss depends on factors such as striga density, host species, land use system, amount of soil nutrients and rainfall pattern (Atera *et al.*, 2012) [11].



Fig 3: A maize field Infested with *Striga hermonthica*

2.3 Genetic resource for *Striga hermonthica* resistance and tolerance in maize

The genetic enhancement of a trait depends on the availability of genetic variability. Sources of resistance to *Striga* have been identified in maize (Amusan, *et al.*, 2008) [89], rice (Gurney *et al.*, 2006) [33], sorghum (Mohamed *et al.*, 2003; Haussmann *et al.*, 2004; Mbuvi *et al.*, 2017) [56, 35, 50] and Cowpea (Menkir, 2006) [52]. Such host-based *Striga* resistance mechanisms act

either before (pre-attachment resistance) or after infection (post-attachment resistance). Potential sources of resistance to *Striga* have been found in a number of heterotic groups of maize (Table 1). Crop wild relatives offer a wide genetic pool for breeding purposes. Although there is little explicit resistance to *Striga* among maize landraces in Africa, some *Striga*-resistant landraces have recently been reported in Kenya.

Table 1: Summary of genetic sources for *Striga* resistance

Germplasm	Current source	References
Wild maize relatives	Lama T. dactyloides- source of Lhf genes for haustorial developmental barriers Z. diploperensis- major source of resistance in maize	(Gurney <i>et al.</i> , 2003) (Amusan <i>et al.</i> , 2008) [89]
Landraces	Sources of horizontal resistance	(Charles <i>et al.</i> (2016)
Inbred lines	IITA and CIMMYT lines	(Menkir, 2006) [52], (Karaya <i>et al.</i> , 2012)
Hybrids	Hybrids Resistant commercial genotypes e.g. Pioneer Hybrids and CGIAR varieties	Chitagu <i>et al.</i> , 2014), (Akinwale <i>et al.</i> 2014)
OPV	IITA populations – e.g. TZL comp1 synw-1 and Acr94TZE Comp s-w	(Menkir and Kling, 2007)

Source: (Admire T.S *et al.*, 2017) [2]

2.4 Screening and evaluation of genetic resources for *Striga* resistance

The development of *Striga*-resistant cultivars has been limited by the lack of reliable screening techniques (Yagoub *et al.*, 2014) [85]. Some of the screening techniques that have been used in the past include laboratory analysis methods, pot screening, and field trials (Rodenburg *et al.*, 2015) [67]. Although the practice of field screening helps in generating statistics about *Striga* infestations under natural conditions, the method is limited by the existing environmental impacts. To circumvent this and initiate reliable screening after attachment, the rhizotron screening system is ideal (Rodenburg *et al.*, 2015) [67]. Rhizotrons are transparent root observation chambers that allow *striga* attached to the host plant to be counted, the phenotype of resistance mechanisms to be assessed, and the effect of *striga* on host biomass to be determined over time and with minimal disruption (Cissoko *et al.*, 2011; Gurney *et al.*, 2006; Rodenburg *et al.*, 2015) [67, 33].

Laboratory screening methods

Traditional breeding methods by screening resistant genotypes under natural or artificial *striga* infestation are very expensive,

time consuming and labor intensive. Using simple laboratory techniques when screening host plants for responses to *Striga* infestation can reveal resistance mechanisms in host plants, increase the efficiency of breeding programs and make it possible to pyramid different resistance mechanisms in a single cultivar for stable resistance responses (Ejeta, 2007) [26]. Laboratory experiments were designed to identify resistance components that together provide the host with general resistance expression during parasite establishment (Ejeta and Gressel 2007) [26]. *In vitro* growth systems allow studying the architecture of host roots and their biochemical resistance mechanisms. Some laboratory assays allow the study of the release of germ inhibitors and haustorial initiation factors as well as hypersensitivity reactions (Ejeta *et al.* 2000) [27]. The paper roll assay was developed to analyze the early stages of *Striga* infection (Ejeta *et al.* 2000) [27]. In this case, preconditioned *striga* seeds are exposed to light and then spread evenly on germination paper moistened with distilled water. The *striga* seeds are then rolled between the germ papers along with lined host seeds. Observations are then made after three weeks when the papers are unrolled to reveal the extent of parasitic attachments to host roots and early resistance mechanisms.

Glasshouse screening

Screening in pots was also an integral part of Striga resistance assessments (Ahonsi *et al.*, 2002). Pots have been used extensively for screening for cultivar resistance, host-parasite nutritional relationships, growth stimulant analysis, and herbicide efficacy. Various pot-strainer techniques such as the polybag and the seed pan have been described in detail by (Rao, 1985). The methods are recommended for their effectiveness in screening for sorghum resistance to *S. hermonthica*. Of note is the development of the Eplee bag pot screening technique developed by (Eplee, 1992). Striga seeds are placed in small micromesh bags, tied to string and buried near the plant roots. At a certain moment, the threads are pulled to observe the germination of the striga. The method can also be used under field conditions to observe Striga germination under natural conditions. Several studies demonstrate the validity of the Eplee-Bag technique as a screening method (Gurney *et al.* 1995; Ahonsi *et al.* 2002). The most important consideration in greenhouse evaluations is its compatibility with experiments on the effectiveness of biological control agents such as *Fusarium oxysporum* f. sp. *striae*. The technique allows for a continuous, uninterrupted assessment of the plant rhizosphere, as demonstrated by (Ahonsi *et al.*, 2002) and (Yonli *et al.* 2006) in their assessment of potential biological control agents in control.

Field techniques

Confusing effects of environmental conditions on polygenic inheritance of traits associated with Striga resistance make field screening imperative despite advances made through laboratory and pot experiments. The art of increasing the accuracy and efficiency of field screening for Striga resistance has been perfected over the years. In corn improvement, an efficient grading scale must be used to estimate the breeding value of a single genotype for Striga. Striga damage rating value, striga occurrence, and agronomic traits that contribute to grain yield are commonly used in selection for resistance in corn. Notwithstanding the shift in focus from selection for tolerance to that in favor of resistance, the Striga damage score still provides a basis for maize improvement for Striga resistance (Menkir 2006) [52]. Tolerance, determined by the Striga damage score, is based on a scale of 19, where 1 means no symptoms of damage and the genotype is considered highly tolerant, while 9 means very susceptible and severe damage (Kim and Adetimirin 1997). It has been found that a low Striga damage rate is directly associated with an increase in grain yield at the same infestation level as the susceptible maize genotypes (Badu-Apraku 2007; Menkir and Kling 2007) [16, 54]. However, selection for tolerance leads to the accumulation of a high striga seed bank. Genetic variants in crop plants have been advanced for further improvement based on their ability to suppress parasitic attachment ladders.

Evaluation of genetic resources for Striga resistance

The development of genetically improved strains with Striga resistance is often straightforward given the availability of reliable sources of resistance and the availability of efficient and easily controllable as well as practical screening methods to create sufficient selection pressure (Rubiales, 2003). Various controlled environment and field screening methods have been developed and applied in Striga improvement programs. Evaluation of germplasm for resistance to parasitic weeds can be performed in controlled and field environments. Controlled environments include laboratory and greenhouse conditions under artificial infestation, while field trials are conducted in

either hotspot areas with additional infestation to increase selective pressure.

Breeding techniques for *S. hermonthica* resistance in maize

Conventional breeding

Conventional breeding techniques have been predominantly used in conferring superior combinations of Striga resistance alleles among susceptible cultivated crops (Menkir *et al.* 2004) [54]. Hybrid breeding, recurrent selection, half-sib selection, full-sib and, S1 family selection schemes have been successfully utilized in developing resistance to most virulent Striga species in legumes and cereal. It is relevant to explore the applicability of most conventional breeding techniques as they have been utilized in various Striga resistance breeding programs. Striga resistance traits have been accumulated successfully through recurrent selection in cereal crops. Through recurrent selection, genetic gains in grain yield in segregating populations has been reported by (Menkir *et al.* 2004) [54] and (Badu-Apraku *et al.* 2006) [15].

The IITA proposes to cluster and mate maize populations of different genetic make-up with contrasting maturity groups and grain colors to create Striga-resistant breeding populations. Quantitative trait loci (QTLs) for resistance to *S. hermonthica* have been identified from local populations including wild relatives and successfully transferred into adaptable maize populations by backcross breeding (Rich and Ejeta 2008). Germplasm obtained by the backcrossing method forms the basis for further development of cultivars to achieve polygenic resistance to *S. hermonthica*. Such inbreds of *Zea diploperennis* and tropical maize were essential for the development of *S. hermonthica*-resistant open-pollinated populations such as *Zea diplo* SYNW-1, TZL Comp SYNW-1. This has been a key resource for municipal corn production systems. Partial resistance to *S. hermonthica* has also been observed in backcross hybrids from a resistant donor *T. dactyloides* (Gurney *et al.* 2003). The backcrossing procedure is straightforward when a starting population or donor with a high frequency of desirable alleles for Striga resistance is available. Rapid progress in establishing resistance to *S. asiatica* can be achieved by identifying a donor with a high dominance of *S. asiatica* resistance genes. In such a scenario, ideal recurring parents would be genotypes that combine early maturity and high yield (Badu-Apraku *et al.* 2006) [15].

Heterosis of hybrid cultivars can be useful to mitigate the effect of *S. hermonthica* on plant production and productivity. With the increasing use of hybrid maize seed in West and Central Africa, IITA was able to accumulate resistance to *S. hermonthica* in hybrid maize by crossing different inbred lines (Menkir *et al.* 2004) [54]. This gives rise to *S. hermonthica*-resistant hybrids that can suppress parasite emergence, with some producing high grain yields at high infestations (Karaya *et al.* 2012). The rapid progress in the development of resistant *S. hermonthica* hybrids in IITA programs can be attributed to the availability of stable resistant genotypes, which have been used as testers to evaluate the broad pool of inbred lines for their overall mating abilities (Menkir *et al.* 2004) [54]. However, conventional breeding has proven to be time-consuming, largely if not entirely dependent on climatic and environmental conditions, and therefore less effective (Ejeta and Gressel, 2007; Rispaill *et al.*, 2007) [26].

Marker assisted breeding for Striga resistance

Although conventional breeding has made significant contributions to improving maize resistance to Striga, it has generally been slowed when targeting the complex quantitative

trait of resistance to Striga. Traditional breeding methods by screening resistant genotypes under natural or artificial striga infestation are very expensive, time consuming and labor intensive. Marker-assisted selection (MAS) is an efficient approach to increase the accuracy and efficiency of selection using markers that are tightly linked to genes to complement phenotypic selection (Srivastava, R.K.; *et al.*, 2020; Wang, X.; *et al.*, 2019) ^[77]. Applying marker-assisted selection (MAS) is a surefire way to improve the efficiency of the Striga resistance breeding program and reduce the inefficiency, lack of precision, and high costs associated with traditional field screening for resistance in host plants. MAS can be facilitated through the use of *in vitro* assays. These laboratory tests can aid in the rapid and effective phenotyping of separating populations for specific resistance mechanisms to Striga. Marker assisted selections are an indispensable element of most breeding programs as they reduce selection errors associated with phenotypic assessments. In addition, phenotyping large pools of germplasm for Striga resistance is expensive, making it challenging to generate sufficient data for high-resolution maker trait association and QTL detection. Currently, there are limited reports on QTL conditioning Striga resistance in maize. However, a recent study involving marker-assisted recurrent selection for grain yield under drought stress and Striga infestation elucidated the importance of this technique in accumulating favorable genes for quantitative traits (Abdulmalik *et al.* 2017).

In general, corn breeding for Striga resistance has relied too heavily on field screening, the accuracy of which can be confounded by a plethora of uncontrolled variation. As reported by (Ejeta and Gressel, 2007) ^[26], some Striga resistance genes are recessive, therefore some sources of resistance can be discarded without molecular mapping. Therefore, the current *S. hermonthica* resistant genetic resource, developed through lengthy and costly breeding cycles, could represent half of its potential. Searching for striga resistance QTL in maize can also use the same approach of testing for resistance QTL for low induction of striga germination and for genomic regions associated with field resistance. This can be achieved by genotyping the maize gene pool at different stages of parasite development. Crosses can be performed between recombinant inbred lines (RIL) derived from susceptible and resistant backgrounds as reported in (Menkir *et al.* 2004) ^[54]. The resulting hybrids can then be examined prior to fingerprinting in agar gel assay tests for histological response mechanisms to parasitic infections for the *lsgs* gene as described by (Haussmann *et al.*, 2004) ^[35].

Role of genetic engineering in improving maize resistance to *S. hermonthica*

Genetic engineering to improve host plant resistance to Striga is a promising approach and offers new opportunities to develop improved cultivars. Over the years, scientists have increased research against parasitic weed species, including Striga, by taking advantage of technological advances that allow insertions, deletions of target genes, manipulation of specific protein sequences, and regulation of plant metabolites involved in the synthesis of cellular structural components (Yoder 2013; Kirigia *et al.* 2014) ^[48]. The studies aimed at the molecular characterization of host-plant-parasite interaction and host resistance through expression analysis of the genes, proteins and metabolites involved in these processes are the subject of increasing interest and offer weed researchers the opportunity to improve the use of genetic transformation tools to control them of parasitic plants (Rispaal *et al.*, 2007; Aly, 2012). So far, no

transgenic Striga-resistant millet plants have been reported and approved for cultivation. However, understanding the biology of the early stages of Striga parasitism will help identify potential barriers to the success of this technology. Although significant strides have been made in genetic transformation, there have been very few reports of sorghum crops lagging behind other cereals such as maize in terms of genetic transformation (Visarada and Kishore, 2015) ^[81]. Regardless of the varying successes recorded by different workers in attempts to induce Striga resistance, all reports reflect a great potential of RNAi in the development of transgenic maize capable of suppressing the parasite's proliferation. (Yoder *et al.* 2009) and (Runo *et al.* 2011) ^[73] have investigated the use of RNA interference (RNAi) technology as a means of enhancing host resistance to parasitic weeds. However, this approach was unsuccessful in controlling the maize striga parasite interaction (Yoder and Scholes, 2010). Interfering double-stranded ribonucleic acid (dsRNA) constructs in maize plants could silence the expression of genes responsible for susceptibility to *S. hermonthica* parasitism (Kirigia *et al.* 2014) ^[48].

Conclusion and Recommendation

Among biotic constraints, *Striga hermonthica* is the most devastating and dominant factor in maize production, causing up to 100% yield loss. The extent of damage depends on the timing and extent of infection. Planting Striga-resistant maize varieties is currently considered the best control strategy and is easy to apply, especially in combination with other management practices. Maize genotypes with significant resistance to *S. hermonthica* are being developed by the International Institute of Tropical Agriculture (IITA) and other national breeding programs. This was a significant milestone in providing resource-poor corn growers in SSA with a cost-effective and effective Striga control option. These genetic resources can serve as useful parents for breeding programs.

Conventional breeding techniques have been predominantly used to confer superior combinations of Striga resistance alleles among susceptible crops. Since resistance to *S. hermonthica* in maize is regulated by many genes, breeding for Striga-resistant cultivars using traditional approaches has been less effective and time-consuming. The identification of QTLs associated with Striga resistance/tolerance would facilitate the rapid development of Striga resistant/tolerant maize genotypes using MAS due to the polygenic nature of the host-parasite relationship and its interaction with environmental factors. Genetic engineering is another option for breeding Striga-resistant strains. It is a very promising approach and offers new opportunities for the development of improved varieties.

Currently, genetic sources of Striga resistance include wild relatives such as *Tripsacum dactyloides*, *Zea diploperennis*, landrace and synthetic species, inbred lines, and OPV. An ideal mix of inbred lines can combine both resistance and tolerance attributes in resulting hybrids. As suggested by many researchers, a successful breeding program is one that can identify and combine striga resistance and tolerance. Therefore, it is essential for any breeding program to provide the breeder with a wide pool of genetic variation. The integration of different breeding techniques to develop *S. hermonthica*-resistant corn varieties would help to reduce losses to corn farmers from this parasite infestation.

References

1. Abdulmalik RO, Menkir A, Meseka SK, Unachukwu N, Ado SG, Olarewaju JD, *et al.* Genetic gains in grain yield of

- a maize population improved through marker assisted recurrent selection under stress and non-stress conditions in West Africa. *Front Plant Sci.* 2017;8:1-11.
2. Admire Tichafa Shayanowako, Mark Laing, Hussein Shimelis. To cite this article: Admire Tichafa Shayanowako, Mark Laing, Hussein Shimelis & Learnmore Mwadzingeni. Resistance breeding and biocontrol of *Striga asiatica* (L.) Kuntze in maize: a review, *Acta Agriculturae Scandinavica, Section B. Soil & Plant Science*; c2017.
 3. Ahonsi MO, Berner DK, Emechebe AM, Lagoke ST. Selection of rhizobacterial strains for suppression of germination of *Striga hermonthica* (Del.) Benth. seeds. *Biol Control.* 2002;24:143–152.
 4. Aigbokhan EI, Berner DK, Musselman LJ, Mignouna HD. Evaluation of variability in *Striga aspera*, *Striga hermonthica* and their hybrids using morphological characters and random amplified polymorphic DNA markers. *Weed Res.* 2000;40:375–386.
 5. Akanvou L, Doku EV, Kling JG. Estimates of genetic variances and interrelationships of traits associated with *Striga* resistance in maize. *Africa Crop Science Journal.* 1997;5:1-8.
 6. Akinwale R, Badu-Apraku B, Fakorede M, Vroh-Bi I. Heterotic grouping of tropical early-maturing maize inbred lines based on combining ability in *Striga*-infested and *Striga*-free environments and the use of SSR markers for genotyping. *Field Crops Res.* 2014;156:48–62.
 7. Aly R. Conventional and biotechnological approaches for control of parasitic weeds. *In vitro Cell Dev Biol.* 2007;43:304-317.
 8. Aly R, Cholakh H, Joel DM, Leibman D, Steinitz B, Zelcer A, *et al.* Gene silencing of mannose 6-phosphate reductase in the parasitic weed *Orobanche aegyptiaca* through the production of homologous dsRNA sequences in the host plant. *Plant Biotechnol J.* 2009;7:487-498.
 9. Amy B Welsh, Kamal I Mohamed. Genetic diversity of *Striga hermonthica* populations in Ethiopia: Evaluating the role of geography and host specificity in shaping population structure. *Int. J Plant Sci.* 2011;172(6):773–782.
 10. Anami S, De Block M, Machuka J, Van Lijsebettens M. Molecular Improvement of Tropical Maize for Drought Stress Tolerance in Sub-Saharan Africa. *Critical Reviews in Plant Sciences.* 2009;28(1-2):16–35.
 11. Atera EA, Itoh K, Azuma T, Ishii T. Farmers' perspectives on the biotic constraint of *Striga hermonthica* and its control in western Kenya. *Weed Biology and Management.* 2012;12(1):53-62.
 12. Atera E, Basin L, Authority D, Atera E. Evaluation of ecologies and severity of *Striga* weed on rice in sub-Saharan Africa; c2014.
 13. Awata LAO, Beyene Y, Gowda M, Suresh LM, Jumbo MB, Tongoona P, *et al.* Genetic analysis of QTL for resistance to Maize Lethal Necrosis in multiple mapping populations. *Genes.* 2020;11:32.
 14. Badu-Apraku B, Akinwale R. Cultivar evaluation and trait analysis of tropical early maturing maize under *Striga*-infested and *Striga*-free environments. *Field Crops Res.* 2011;121:186–194.
 15. Badu-Apraku B, Menkir A, Fakorede M, Lum AF, Obeng-Antwi K. Multivariate analyses of the genetic diversity of fortyseven *Striga* resistant tropical early maturing maize inbred lines. *Maydica.* 2006;51:551.
 16. Badu-Apraku B, Fontem Lum A, Fakorede MAB, Menkir A, Chabi Y, The C, *et al.* Performance of early maize cultivars derived from recurrent selection for grain yield and *Striga* resistance. *Crop Sci.* (In press); c2007.
 17. Badu-Apraku B, Fakorede MAB, Oyekunle M, Yallou GC, Obeng-Antwi K, Haruna A, *et al.* Gains in grain yield of early maize cultivars developed during three breeding eras under multiple environments. *Crop Science.* 2015;55:527-539.
 18. Badu-Apraku B, Fakorede M, Gedil M, Annor B, Talabi AO, Akaogu IC, *et al.* Heterotic patterns of IITA and CIMMYT early-maturing yellow maize inbreds under contrasting environments. *Agronomy Journal.* 2016;108:1321-1336.
 19. Badu-Apraku B, Fakorede MAB. *Advances in Genetic Enhancement of Early and Extra-Early Maize for Sub-Saharan Africa*; Springer: Cham, Switzerland; c2017.
 20. Berner D, Winslow M, Awad A, Cardwell K, Raj DM, Kim S. *Striga* research methods. Manual, the Pan-African *Striga* Control Network (PASCON) and the International Institute of Tropical Agriculture. Ibadan, Nigeria: International Institute of Tropical Agriculture (IITA); c1997.
 21. Bharathalakshmi, Werth CR, Musselman LJ. A study of genetic diversity among host-specific populations of the witch weed *Striga hermonthica* (Scrophulariaceae) in Africa. *Plant Syst Evol* 172: *Biotechnol Dev Monitor.* 1990;448(48):6–7.
 22. Bouwmeester HJ, Matusova R, Zhongkui S, Beale MH. Secondary metabolite signalling in host–parasitic plant interactions. *Current Opinion in Plant Biology.* 2003;6(4):358-364.
 23. Chitagu M, Rugare JT, Mabasa S. Screening maize (*Zea mays*) genotypes for tolerance to witchweed (*Striga asiatica* L. Kuntze) Infection. *J Agr Sci.* 2014;6:160–169.
 24. De Groote H. *Striga* economics. Integrating new technologies for *Striga* control: Towards ending the witch-hunt. Singapore: World Scientific Publishing Company; c2007.
 25. Diallo A, Kanampiu F, Mugo S, De Groote H, Mbogo P. *Herbicide Resistant Maize: A Novel Method to Control Striga in Africa.* Paper presented at the 5th West and Central Africa Biennial Regional Maize Workshop, IITA-Cotonou, Benin Republic; c2005.
 26. Ejeta Gebisa, Gressel J. Integrating new technologies for *Striga* control: Towards ending the witch-hunt. World Scientific; c2007.
 27. Ejeta Gebisa, Mohammed A, Rich P, Melake-Berhan A, Housley TL, Hess DE. Selection for specific mechanisms of resistance to *Striga* in sorghum. *Breeding for Striga resistance in cereals*; c2000.
 28. Ejeta Gebisa. *Breeding for resistance in sorghum: Exploitation of an intricate host–parasite biology.* *Crop Sci.* 2007;47:S-216–S227.
 29. Eplee RE. *Witchweed (Striga asiatica): an overview of management strategies in the USA.* *Crop Prot.* 1992;11:3–7.
 30. FAO. *Sowing the Seeds of Peace for Food Security: Disentangling the nexus between conflict, food security and peace*, edited by C. Holleman, J. Jackson, M.V. Sánchez & R. Vos. FAO Agricultural Development Economics Technical Study 2. Rome; c2017.
 31. Gasura E, Setimela P, Mabasa S, Rwafa R, Kageler S, Nyakurwa C. Response of IITA maize inbred lines bred for *Striga hermonthica* resistance to *Striga asiatica* and associated resistance mechanisms in southern Africa. *Euphytica* 2019, 215, 151. [CrossRef] *Agronomy.* 2020;10:1168 16 of 18

32. Gressel J, Hanafi A, Head G, Marasas W, Obilana AB, Ochanda J, *et al.* Major heretofore intractable biotic constraints to African food security that may be amenable to novel biotechnological solutions. *Crop Protection*. 2004;23(8):661–689.
33. Gurney AL, Slate J, Press MC, Scholes JD. A novel form of resistance in rice to the angiosperm parasite *Striga hermonthica*. *New Phytologist*. 2006;169(1):199-208.
34. Gutierrez-Marcos JF, Pennington PD, Costa LM, Dickinson HG. Imprinting in the endosperm: A possible role in preventing wide hybridization. *Philos Trans R Soc Lond*. 2003;358:1105-1111.
35. Haussmann B, Hess D, Omany G, Folkertsma R, Reddy B, Kayentao M, *et al.* Genomic regions influencing resistance to the parasitic weed *Striga hermonthica* in two recombinant inbred populations of sorghum. *Theor Appl Genet*. 2004;109:1005-1016.
36. Haussmann B, Hess D, Reddy B, Mukuru S, Kayentao M, Welz H, *et al.* Pattern analysis of genotype× environment interaction for *Striga* resistance and grain yield in African sorghum trials. *Euphytica*. 2001;122:297-308.
37. Haussmann BI, Hess DE, Welz H-G, Geiger HH. Improved methodologies for breeding *Striga*-resistant sorghums. *Field Crops Res*. 2000;66:195-211.
38. Haussmann BIG, Hess DE, Omany GO, Folkertsma RT, Reddy BVS, Kayentao M, *et al.* Genomic regions influencing resistance to the parasitic weed *Striga hermonthica* in two recombinant inbred populations of sorghum. *Theoretical and Applied Genetics*. 2004;109:1005-1016.
39. Irving LJ, Cameron DD. You are what you eat: interactions between root parasitic plants and their hosts. *Adv. Bot. Res*. 2009;50:87-138.
40. Joel DM, Hershenthorn J, Eizenberg H, Aly R, Ejeta G, Rich PJ, *et al.* Biology and management of weedy root parasites. *Hort. Rev*. 2007;33:267-349.
41. Kanampiu F, Ransom J, Gressel J, Jewell D, Friesen D, Grimanelli D, *et al.* Appropriateness of biotechnology to African agriculture: *Striga* and maize as paradigms. *Plant Cell Tiss Org*. 2002;69:105-110.
42. Kanampiu FK, Kabambe V, Massawe C, Jasi L, Friesen D, Ransom JK, *et al.* Multi-site, multi-season field tests demonstrate that herbicide seed-coating herbicide-resistance maize controls *Striga* spp. and increases yields in several African countries. *Crop Prot*. 2003;22:697-706.
43. Kebede Mulugeta, Dorothee Ehrlich, Pierre Taberlet, Nemomissa Sileshi, Christian Brochmann. Phylogeography and conservation genetics of a giant lobelia (*Lobelia giberroa*) in Ethiopian and tropical East African mountains. *Mol Ecol*. 2007;16:1233-1243.
44. Khan ZR, Hassanali A, Overholt W, Khamis TM, Hooper AM, Pickett JA, *et al.* Combined control of *Striga hermonthica* and stem borers by maize–*Desmodium* spp. intercrops. *Crop Protection*. 2006;25(9):989-995.
45. Kim SK, Adetimirin VO. *Striga hermonthica* seed inoculum rate effects on maize hybrid tolerance and susceptibility expression. *Crop Sci*. 1997;37:1066-1071.
46. Kim SK. Genetics of maize tolerance of *Striga hermonthica*. *Crop Science*. 1994;34(4):900-907.
47. Kingdon J. *Island Africa: The evolution of Africa's rare animals and plants*. Collins, London; c1990.
48. Kirigia D, Runo S, Alakonya A. A virus-induced gene silencing (VIGS) system for functional genomics in the parasitic plant *Striga hermonthica*. *Plant Methods*. 2014;10:16.
49. Mather K, Jinks JL. *Biometrical Genetics* (3rd ed.). London, UK: Chapman and Hall; c1982.
50. Mbuvi DA, Masiga CW, Kuria E, Masanga J, Wamalwa M, Mohamed A, *et al.* Novel Sources of Witch weed (*Striga*) Resistance from Wild Sorghum Accessions. *Frontiers in Plant Science*. 2017;8:116.
51. Melaku Gedil M, Abebe Menkir A. An integrated molecular and conventional breeding scheme for enhancing genetic gain in maize in Africa. *Front. Plant Sci*. 2019;10:1430.
52. Menkir A. Assessment of reactions of diverse maize inbred lines to *Strigahermonthica* (Del.) Benth. *Plant Breeding*. 2006;125:131-139.
53. Menkir A, Chikoye D, Lum F. Incorporating an herbicide resistance gene into tropical maize with inherent polygenic resistance to control *Striga hermonthica* (Del.) Benth. *Plant Breeding*. 2010;129:385–392.
54. Menkir A, Kling JG, Badu-Apraku B. Recent advances in breeding maize for resistance to *Striga hermonthica* (del.) Benth; c2004.
55. Mohamed AH, Housley T, Ejeta G. An *in vitro* technique for studying specific *Striga* resistance mechanisms in sorghum. *African J Agr Res*. 2010;5:1868–1875.
56. Mohamed A, Ellicott A, Housley TL, Ejeta G. Hypersensitive response to *Striga* infection in Sorghum. *Crop Sci*. 43, 1320 1324. *Molecular Plant Pathology*. 2003;14(9):861–869.
57. Mutengwa C, Tongoona P, Sithole-Niang I. Genetic studies and a search for molecular markers that are linked to *Striga asiatica* resistance in sorghum. *Aft J Biotechnol*. 2005;4:1355-1361.
58. Mwangi P, Ely A. Assessing risks and benefits: Bt maize in Kenya; c2001.
59. Obilana AT. Inheritance of resistance to *Striga* (*Striga hermonthica* Benth.) in sorghum. *Prot. Ecol*. 1984;7:305-311.
60. Olivier A, Glaszmann JC, Lanaud C, Leroux GD. Population structure, genetic diversity and host specificity of the parasitic weed *Striga hermonthica* (Scrophulariaceae) in Sahel. *Plant Syst Evol*. 1998;209:33-45.
61. Pathi KM, Tula S, Huda KMK, Srivastava VK, Tuteja N. An efficient and rapid regeneration via multiple shoot induction from mature seed derived embryogenic and organogenic callus of Indian maize (*Zea mays* L.). *Plant Signaling & Behavior*. 2013;18(3):303-315.
62. Ransom J, Kanampiu F, Gressel J, De Groot H, Burnet M, Odhiambo G. Herbicide applied to imidazolinone resistant-maize seed as a *Striga* control option for smallscale African farmers. *Weed Sci*. 2012;60:283-289.
63. Ranum P, Peña-Rosas JP, Garcia-Casal MN. Global maize production, utilization, and consumption. *Ann NY Acad Sci*. 2014;1312:105-112.
64. Reda F, Butler L, Ejeta G, Ransom K. Screening of maize genotypes for low *Striga asiatica* stimulant production using the agar gel technique'. *Afri Crop Sci J*. 1994;2:173-177.
65. Redinbaugh M, Pratt R. Virus resistance. In *Handbook of Maize: It's Biology*; Bennetzen JL, Hake SC, Eds; Springer: New York, NY, USA; c2009. p. 251-268.
66. Reynolds TW, Waddington SR, Anderson CL, Chew A, True Z, Cullen A. Environmental impacts and constraints associated with the production of major food crops in Sub-Saharan Africa and South Asia. *Food Security*. 2015;7(4):795-822.

67. Rodenburg J, Bastiaans L. Host-plant defence against *Striga* spp.: reconsidering the role of tolerance. *Weed Res.* 2015;51:438-441.
68. Rodenburg J, Bastiaans L, Kropff MJ, van Ast A. Effects of host plant genotype and seed bank density on *Striga* reproduction. *Weed Res.* 2006;46:251-263.
69. Rouf Shah T, Prasad K, Kumar P. A potential source of human nutrition and health: A review. *Cogent Food & Agriculture.* 2016;2(1):1-9.
70. Rubiales D, Fernández-Aparicio M, Pérez-de-Luque A, Castillejo MA, Prats E, Sillero JC, *et al.* Breeding approaches for crenate broomrape (*Orobanche crenata* Forsk.) management in pea (*Pisum sativum* L.). *Pest Manag Sci.* 2009;65:553-559.
71. Rubiales D, Pérez-de-Luque A, Fernández-Aparicio M, Sillero JC, Román B, Kharrat M, *et al.* Screening techniques and sources of resistance against parasitic weeds in grain legumes. *Euphytica.* 2006;147:187-199.
72. Rubiales D. Parasitic plants, wild relatives and the nature of resistance. *New Phytol.* 2003;160:459.
73. Runo S, Alakonya A, Machuka J, Sinha N. RNA interference as a resistance mechanism against crop parasites in Africa: A 'Trojan horse' approach. *Pest Manag Sci.* 2011;67:129-136.
74. Ruyter-Spira C, Kohlen W, Charnikhova T, van Zeijl A, van Bezouwen L, de Ruijter N, *et al.* Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in *Arabidopsis*: another belowground role for strigolactones? *Plant Physiology.* 2011;155(2):721-34.
75. Safa SB, BM Jones, LJ Musselman. Mechanisms favoring outbreeding in *Striga hermonthica* (Scrophulariaceae). *New Phytol.* 1984;96:299-305.
76. Spallek T, Mutuku M, Shirasu K. The genus *Striga*: A witch profile; c2013.
77. Srivastava RK, Singh RB, Pujarula VL, Bollam S, Pusuluri M, Chellapilla TS, *et al.* Genome-wide association studies and genomic selection in PearlMillet: Advances and prospects. *Front. Genet.* 2020;10:1389.
78. Teka HB. African Journal of Plant Science Advance research on *Striga* control: A review. 2014;8(11):492-506.
79. Thomson JA. The role of biotechnology for agricultural sustainability in Africa. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 2008;363(1492):905-913.
80. Van Bueren ETL, Backes G, De Vriend H, Østergård H. The role of molecular markers and marker assisted selection in breeding for organic agriculture. *Euphytica.* 2010;175:51-64.
81. Visarada KBRS, Kishore NS. Advances in Genetic Transformation. In: Madhusudhana, R., *et al.* (Eds.), *Sorghum Molecular Breeding.* Springer (India) Pvt. Ltd.; c2015. p. 199-215.
82. Wang X, Liu H, Pang M, Fu B, Yu X, He S, *et al.* Construction of a high-density genetic linkage map and mapping of quantitative trait loci for growth-related traits in silver carp (*Hypophthalmichthys molitrix*). *Sci Rep.* 2019;9:17506.
83. Wondimu Bayu, Selomon Binor, Legesse Admasu. Tolerance of sorghum landraces and varieties to *Striga* (*Striga hermonthica*) infestation in Ethiopia. *Acta Agronomica Hungarica.* 2001;49(4):343-349.
84. Woodcock CM. Control of witchweed *Striga hermonthica* by intercropping with *Desmodium* spp., and the mechanism defined as allelopathic. *Journal of Chemical Ecology.* 2002;28(9):1871-1885.
85. Yagoub SO, Hassan MM, Gani MEA, El A, Babiker GE. Screening Sorghum for Resistance to *Striga hermonthica* (Del.) Benth. 2014;3(3):1-5.
86. Yoder JI. Host-plant recognition by parasitic Scrophulariaceae. *Current Opinion in Plant Biology.* 2001;4(4):359-365.
87. Yoder JI, Gunathilake P, Wu B, Tomilova N, Tomilov AA. Engineering host resistance against parasitic weeds with RNA interference. *Pest Manage. Sci.* 2009;65:460-466.
88. Yoder JI, Scholes JD. Host plant resistance to parasitic weeds; recent progress and bottlenecks. *Curr; c2010.*
89. Amusan OA, Massengill LW, Baze MP, Sternberg AL, Witulski AF, Bhuva BL, Black JD. Single event upsets in deep-submicrometer technologies due to charge sharing. *IEEE Transactions on Device and Materials Reliability.* 2008;8(3):582-9.
90. Keyes CL. Promoting and protecting mental health as flourishing: a complementary strategy for improving national mental health. *American psychologist.* 2007;62(2):95.