



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2024; SP-7(11): 395-399

Received: 01-08-2024

Accepted: 09-09-2024

**SP Bhagat**

M.Sc. Scholar, Department of  
Agronomy, PGI, MPKV, Rahuri,  
Maharashtra, India

**NK Medhe**

Cotton Agronomist, Cotton  
Improvement Project, MPKV,  
Rahuri, Maharashtra, India

**SS Ilhe**

Professor, Department of  
Agronomy, PGI, MPKV, Rahuri,  
Maharashtra, India

**NJ Danawale**

Chief Scientist (Seeds), Associate  
Professor, Department of  
Agronomy, PGI, MPKV, Rahuri,  
Maharashtra, India

**NK Bhute**

Cotton Entomologist, Cotton  
Improvement Project, MPKV,  
Rahuri, Maharashtra, India

**PS Bodake**

Head, Department of Agronomy,  
PGI, MPKV, Rahuri,  
Maharashtra, India

**MR Patil**

Associate Professor, Department of  
Agricultural Statistics, PGI,  
MPKV, Rahuri, Maharashtra,  
India

**Corresponding Author:**

**SP Bhagat**

M.Sc. Scholar, Department of  
Agronomy, PGI, MPKV, Rahuri,  
Maharashtra, India

## Yield maximization of *Bt.* cotton by integrated crop management techniques

**SP Bhagat, NK Medhe, SS Ilhe, NJ Danawale, NK Bhute, PS Bodake and MR Patil**

**DOI:** <https://doi.org/10.33545/2618060X.2024.v7.i11Sf.2019>

### Abstract

A field experiment entitled, “Yield maximization of *Bt.* cotton by integrated crop management techniques” was carried out during *kharif*, 2023 at Cotton Improvement Project, Mahatma Phule Krishi Vidyapeeth, Rahuri, Maharashtra (India). The experiment was laid out in Randomized Block Design (RBD) with three replications and nine treatments. The results of present investigation revealed that, yield and yield contributing characters of *Bt.* cotton were significantly influenced by different treatments. Treatment T<sub>1</sub> i.e., wider spacing (90 cm × 90 cm) recorded significantly maximum lint weight boll<sup>-1</sup> (1.36 g), seed weight boll<sup>-1</sup> (2.64 g), boll weight (4.00 g), number of green (unopened) bolls plant<sup>-1</sup> (2.60), number of picked bolls plant<sup>-1</sup> (45.09) and seed cotton yield plant<sup>-1</sup> (180.36 g). The highest seed cotton yield (4162 kg ha<sup>-1</sup>) and stalk yield (6202 kg ha<sup>-1</sup>) were recorded with treatment T<sub>9</sub> (Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS). The quality parameters viz., ginning percentage (%), lint index (%), micronaire value (µg inch<sup>-1</sup>), tensile strength (g tex<sup>-1</sup>), 2.5% span length (mm) and uniformity ratio (%) were not influenced significantly due to different treatments. Treatment T<sub>9</sub> (Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) recorded highest cost of cultivation (₹ 1,27,328), gross monetary returns (₹ 2,81,780), net monetary returns (₹ 1,54,452) and B:C ratio (2.21) than rest of all treatments.

**Keywords:** *Bt.* cotton, closer spacing, mepiquat chloride, detopping, pruning of monopodia

### Introduction

Cotton is one of the predominant fiber crop and plays a pivotal role in agriculture, industrial development, employment generation and economic development of India. It is called as "King of fibers" and "White gold" due to higher economical values among all cash crops in India. Nearly one third of India's export earnings are from the textile sectors. Apart from its fiber, it contributes considerably to the non-edible oil pool of our country and likely to play pivotal role in paper, particle board and card board industries, etc. in coming decades (Elamathy *et al.*, 2022) [5].

Cotton is a sub-tropical plant having indeterminate growth habit. Among various agronomic practices to boost up crop productivity, plant population is one of the most important factor for efficient utilization of available resources. The manipulation of plant density and crop geometry is a time-tested agronomic technique for achieving higher crop yield. In wider spacing, yield reduces due to less plant per unit area and closer spacing reduces yield due to competition within the plants (Gohil *et al.*, 2023) [6].

Managing the equilibrium between vegetative and reproductive growth is an important part in cotton production. This can be achieved by detopping and pruning or removal of monopodial branches. Detopping is the removal of top terminal portion after prominent vegetative growth stage. This may encourage growth of already formed sympodia as well as formation and development of fruiting bodies. Detopping may also help to reduce sucking pests and bollworm infestation, as it avoids fresh growth emergence which is unwanted at particular stage (Kakde *et al.*, 2023) [7].

In India, cotton is grown under diverse agro-climatic conditions. With the better understanding of plant growth habits, new agricultural practices are developed for higher productivity of the crops. Production and productivity of cotton crop increased with the introduction of genetically modified *Bt* cotton in the country. For fully exploitation of the crop yield, there is a need to develop new crop management practices.

### Materials and Methods

The field experiment entitled, “Yield maximization of *Bt* cotton by integrated crop management techniques” was conducted at Cotton Improvement Project, Mahatma Phule Krishi Vidyapeeth, Rahuri during *kharif* season, 2023-2024. The soil of the experimental plot was clayey in texture, low in available nitrogen (177.20 kg ha<sup>-1</sup>), medium in available phosphorus (26.50 kg ha<sup>-1</sup>), very high in available potassium (680 kg ha<sup>-1</sup>) and slightly alkaline in reaction (pH 7.7). The EC and organic carbon content was 0.38 dSm<sup>-1</sup> and 0.57% respectively. The experiment was laid out in Randomized Block Design (RBD) with three replications and nine treatments viz., T<sub>1</sub>: Farmer’s practice (90 cm × 90 cm spacing), T<sub>2</sub>: Cotton in closer spacing

(90 cm × 30 cm spacing), T<sub>3</sub>: Cotton in closer spacing + Detopping at 100 cm height, T<sub>4</sub>: Cotton in closer spacing + Pruning of monopodia at 45 DAS, T<sub>5</sub>: Cotton in closer spacing + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS, T<sub>6</sub>: Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS, T<sub>7</sub>: Cotton in closer spacing + Detopping at 100 cm height + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS, T<sub>8</sub>: Cotton in closer spacing + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS, T<sub>9</sub>: Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS. Seeds of cotton variety Ajeet-155 were sown in line by dibbling method and two hand weedings were given to keep the plots weed free. The recommended fertilizer dose (125:65:65 N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O kg ha<sup>-1</sup>) was applied through urea, SSP and MOP along with FYM.

### Results and Discussion

#### Yield Contributing Characters Studies

**Table 1:** Number of picked bolls plant<sup>-1</sup> and boll weight as influenced by integrated crop management techniques

| Treatment details |                                                                                                                                                       | Number of picked bolls plant <sup>-1</sup> | Boll weight (g) |
|-------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------|-----------------|
| T <sub>1</sub>    | Farmer’s practice (90 cm × 90 cm spacing)                                                                                                             | 45.09                                      | 4.00            |
| T <sub>2</sub>    | Cotton in closer spacing (90 cm × 30 cm spacing)                                                                                                      | 24.58                                      | 3.11            |
| T <sub>3</sub>    | Cotton in closer spacing + Detopping at 100 cm height                                                                                                 | 27.01                                      | 3.38            |
| T <sub>4</sub>    | Cotton in closer spacing + Pruning of monopodia at 45 DAS                                                                                             | 27.15                                      | 3.43            |
| T <sub>5</sub>    | Cotton in closer spacing + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                                               | 27.41                                      | 3.54            |
| T <sub>6</sub>    | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS                                                                | 29.21                                      | 3.70            |
| T <sub>7</sub>    | Cotton in closer spacing + Detopping at 100 cm height + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                  | 27.76                                      | 3.64            |
| T <sub>8</sub>    | Cotton in closer spacing + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                              | 28.16                                      | 3.66            |
| T <sub>9</sub>    | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS | 29.23                                      | 3.86            |
|                   | S.E m ±                                                                                                                                               | 5.05                                       | 0.10            |
|                   | C.D. at 0.05                                                                                                                                          | 16.63                                      | 0.32            |
|                   | General mean                                                                                                                                          | 29.51                                      | 3.59            |

The statistically analysed data revealed that yield and yield contributing characters of *Bt* cotton at harvest was significantly influenced due to various treatments. Farmer’s practice (90 cm × 90 cm spacing) recorded significantly higher number of picked bolls plant<sup>-1</sup> at harvest, however, it was at par with treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) and treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS). Lower

number of picked bolls plant<sup>-1</sup> was recorded at treatment T<sub>2</sub> i.e., Cotton in closer spacing (90 cm × 30 cm). More number of picked bolls plant<sup>-1</sup> in farmer’s practice is might be due to the substantial space available for growth, more photosynthetic efficiency, frequent availability of water and nutrients, which resulted in increase in fruiting points, fruiting period, fruit retention and ultimately greater picked bolls plant<sup>-1</sup>. Similar results were obtained by Munir *et al.* (2015) [10] and Uma Maheswari *et al.* (2019) [16].

**Table 2:** Seed cotton yield plant<sup>-1</sup>, seed cotton yield and stalk yield as influenced by integrated crop management techniques

| Treatment details |                                                                                                                      | Seed cotton yield plant <sup>-1</sup> (g) | Seed cotton yield (kg ha <sup>-1</sup> ) | Stalk yield (kg ha <sup>-1</sup> ) |
|-------------------|----------------------------------------------------------------------------------------------------------------------|-------------------------------------------|------------------------------------------|------------------------------------|
| T <sub>1</sub>    | Farmer’s practice (90 cm × 90 cm spacing)                                                                            | 180.36                                    | 2227                                     | 4231                               |
| T <sub>2</sub>    | Cotton in closer spacing (90 cm × 30 cm spacing)                                                                     | 76.43                                     | 2621                                     | 4980                               |
| T <sub>3</sub>    | Cotton in closer spacing + Detopping at 100 cm height                                                                | 91.30                                     | 3157                                     | 5367                               |
| T <sub>4</sub>    | Cotton in closer spacing + Pruning of monopodia at 45 DAS                                                            | 93.13                                     | 3258                                     | 5539                               |
| T <sub>5</sub>    | Cotton in closer spacing + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                              | 97.01                                     | 3432                                     | 5835                               |
| T <sub>6</sub>    | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS                               | 108.08                                    | 4012                                     | 6178                               |
| T <sub>7</sub>    | Cotton in closer spacing + Detopping at 100 cm height + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS | 101.02                                    | 3589                                     | 5922                               |
| T <sub>8</sub>    | Cotton in closer spacing + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g                   | 103.05                                    | 3703                                     | 6036                               |

|                | a.i. at 45 and 60 DAS                                                                                                                                 |        |         |         |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|--------|---------|---------|
| T <sub>9</sub> | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS | 112.80 | 4162    | 6202    |
|                | S.E m +                                                                                                                                               | 24.40  | 74.49   | 44.13   |
|                | C.D. at 0.05                                                                                                                                          | 73.20  | 223.32  | 132.39  |
|                | General mean                                                                                                                                          | 107.02 | 3351.28 | 5587.73 |

Farmer's practice (90 cm × 90 cm spacing) recorded significantly higher boll weight at harvest, however, it was at par with treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) and treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS). Lower boll weight was recorded at treatment T<sub>2</sub> i.e., closer spacing (90 cm × 30 cm). Maximum boll weight in wider spacing is might be due to improved source-sink relationship and better partitioning of photoassimilates into reproductive structures as reported by Kaul *et al.* (2016) [8] and Uma Maheswari *et al.* (2019) [16].

Farmer's practice (90 cm × 90 cm spacing) recorded significantly higher seed cotton yield plant<sup>-1</sup> at harvest, however, it was at par with treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) and treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS). Lower seed cotton yield plant<sup>-1</sup> was recorded at treatment T<sub>2</sub> i.e., closer spacing (90 cm × 30 cm). Under wider spacing availability of photosynthates to individual plant was more to produce maximum seed cotton yield plant<sup>-1</sup> as compared to closer plant spacing. This might be due to overall improvement in growth attributes and its positive effect on number of bolls plant<sup>-1</sup> and seed cotton weight boll<sup>-1</sup> under wider plant spacing. The above results are in conformity with the findings of Khetre *et al.* (2018) [9] and Gohil *et al.* (2023) [6].

Treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) recorded significantly higher seed cotton yield at harvest, however, it was at par with treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height +

Pruning of monopodia at 45 DAS). Lower seed cotton yield was recorded at farmer's practice (90 cm × 90 cm). The remarkable improvement in seed cotton yield in treatment combination of integrated crop management techniques could be due better partitioning of photo assimilates towards reproductive structures, improved source-sink relationship and better translocation of metabolites towards growing and reproductive sink due to retardation of excessive vegetative growth. The results are in conformity with those obtained earlier by Bhalariao and Gaikwad (2010) [1], Shekar *et al.* (2015) [13], Parihar *et al.* (2018) [11] and Uma Maheswari *et al.* (2019) [16].

Treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) recorded significantly higher stalk yield at harvest, however, it was at par with treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS). Lower stalk yield was recorded at farmer's practice (90 cm × 90 cm). The higher cotton stalk yield under closer spacing is mainly due to higher plant population as compared to the wider spacing. These findings are in line with those supported by Bhalariao and Gaikwad (2010) [1], Shekar *et al.* (2015) [13] and Khetre *et al.* (2018) [9].

### Fiber Quality Studies

The statistically analysed data revealed that the quality parameters of *Bt.* cotton viz., ginning percentage (%), lint index (%), micronaire value (µg inch<sup>-1</sup>), tensile strength (g tex<sup>-1</sup>), 2.5% span length (mm) and uniformity ratio (%) were not influenced significantly due to different treatments. Similar results were observed by Dhillon *et al.* (2006) [4], Pendharkar *et al.* (2010) [12], Dadgale *et al.* (2012) [3], Kaul *et al.* (2016) [8] and Chetana *et al.* (2023) [2].

**Table 3:** Quality parameters as influenced by integrated crop management techniques

|                | Treatment details                                                                                                                                     | Ginning percent (%) | Lint index (%) | Micronaire value (µg inch <sup>-1</sup> ) | Tensile strength (g tex <sup>-1</sup> ) | 2.5% span length (mm) | Uniformity ratio (%) |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|----------------|-------------------------------------------|-----------------------------------------|-----------------------|----------------------|
| T <sub>1</sub> | Farmer's practice (90 cm × 90 cm spacing)                                                                                                             | 34.01               | 4.02           | 3.62                                      | 27.93                                   | 26.90                 | 83.67                |
| T <sub>2</sub> | Cotton in closer spacing (90 cm × 30 cm spacing)                                                                                                      | 34.08               | 4.07           | 3.61                                      | 28.07                                   | 26.80                 | 85.67                |
| T <sub>3</sub> | Cotton in closer spacing + Detopping at 100 cm height                                                                                                 | 34.23               | 4.05           | 3.50                                      | 27.90                                   | 27.07                 | 84.00                |
| T <sub>4</sub> | Cotton in closer spacing + Pruning of monopodia at 45 DAS                                                                                             | 34.17               | 4.12           | 3.73                                      | 29.20                                   | 27.17                 | 85.00                |
| T <sub>5</sub> | Cotton in closer spacing + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                                               | 33.55               | 4.16           | 3.61                                      | 28.70                                   | 27.57                 | 84.00                |
| T <sub>6</sub> | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS                                                                | 34.57               | 4.26           | 3.63                                      | 27.40                                   | 26.00                 | 83.00                |
| T <sub>7</sub> | Cotton in closer spacing + Detopping at 100 cm height + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                  | 34.67               | 4.16           | 3.69                                      | 29.00                                   | 27.60                 | 84.00                |
| T <sub>8</sub> | Cotton in closer spacing + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                              | 34.73               | 4.23           | 3.50                                      | 27.93                                   | 27.30                 | 85.00                |
| T <sub>9</sub> | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS | 34.80               | 4.39           | 3.70                                      | 29.60                                   | 27.20                 | 83.00                |
|                | S.E m +                                                                                                                                               | 0.26                | 0.11           | 0.11                                      | 0.47                                    | 0.32                  | 1.67                 |
|                | C.D. at 0.05                                                                                                                                          | NS                  | NS             | NS                                        | NS                                      | NS                    | NS                   |
|                | General mean                                                                                                                                          | 34.31               | 4.16           | 3.62                                      | 28.41                                   | 27.07                 | 84.15                |

## Economic Studies

**Table 4:** Economic studies of *Bt.* cotton as influenced by integrated crop management techniques

|                | Treatment details                                                                                                                                     | Cost of cultivation<br>(₹ ha <sup>-1</sup> ) | Gross monetary<br>returns (₹ ha <sup>-1</sup> ) | Net monetary<br>returns (₹ ha <sup>-1</sup> ) | B:C<br>ratio |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|-------------------------------------------------|-----------------------------------------------|--------------|
| T <sub>1</sub> | Farmer's practice (90 cm × 90 cm spacing)                                                                                                             | 90567                                        | 150748                                          | 60181                                         | 1.66         |
| T <sub>2</sub> | Cotton in closer spacing (90 cm × 30 cm spacing)                                                                                                      | 100143                                       | 177463                                          | 77319                                         | 1.77         |
| T <sub>3</sub> | Cotton in closer spacing + Detopping at 100 cm height                                                                                                 | 109060                                       | 213741                                          | 104681                                        | 1.96         |
| T <sub>4</sub> | Cotton in closer spacing + Pruning of monopodia at 45 DAS                                                                                             | 111699                                       | 220577                                          | 108878                                        | 1.97         |
| T <sub>5</sub> | Cotton in closer spacing + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                                               | 111837                                       | 232365                                          | 120528                                        | 2.07         |
| T <sub>6</sub> | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS                                                                | 122993                                       | 271593                                          | 148601                                        | 2.20         |
| T <sub>7</sub> | Cotton in closer spacing + Detopping at 100 cm height + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                                  | 116895                                       | 242970                                          | 126074                                        | 2.08         |
| T <sub>8</sub> | Cotton in closer spacing + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS                              | 119535                                       | 250699                                          | 131164                                        | 2.10         |
| T <sub>9</sub> | Cotton in closer spacing + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS | 127328                                       | 281780                                          | 154452                                        | 2.21         |
|                | S.E m +                                                                                                                                               | -                                            | 3800                                            | 3800                                          | -            |
|                | C.D. at 0.05                                                                                                                                          | -                                            | 11394                                           | 11394                                         | -            |
|                | General mean                                                                                                                                          | 112228                                       | 226881                                          | 114653                                        | 2.00         |

In economic studies, treatment T<sub>9</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS + Two sprays of Mepiquat chloride @ 25 g a.i. at 45 and 60 DAS) recorded significantly higher cost of cultivation, gross monetary returns, net monetary returns and B:C ratio. However, it was at par with treatment T<sub>6</sub> (Cotton in closer spacing (90 cm × 30 cm) + Detopping at 100 cm height + Pruning of monopodia at 45 DAS) in case of GMR, NMR and B:C ratio. Lower economic values were recorded at farmer's practice (90 cm × 90 cm). This was mainly due to yield obtained in closer spacing was higher than that of wider spacing. Similar results were observed by Shwetha *et al.* (2009)<sup>[14]</sup>, Bhalerao and Gaikwad (2010)<sup>[1]</sup>, Dadgale *et al.* (2012)<sup>[3]</sup>, Singh *et al.* (2017)<sup>[15]</sup>, Khetre *et al.* (2018)<sup>[9]</sup> and Chetana *et al.* (2023)<sup>[2]</sup>.

## Conclusion

On the basis of present investigation, it can be concluded that the *Kharif* season *Bt.* Cotton shall be sown at 90 cm x 30 cm spacing along with complete package of integrated crop management techniques including detopping at 100 cm height, pruning of monopodia at 45 DAS and application of mepiquat chloride @ 25 g a.i. at 45 and 60 DAS for getting maximum yields, gross monetary returns, net monetary returns and benefit cost ratio in medium black soil of Maharashtra.

By keeping in view the hazardous effects of application of chemicals such as mepiquat chloride on soil and environment, it is recommended to practice closer spacing at 90 cm x 30 cm along with integrated crop management techniques including detopping at 100 cm height and pruning of monopodia at 45 DAS for sustainable production and productivity in *Bt.* cotton.

## Acknowledgement

We are very thankful to the Post Graduate Institute and Cotton Improvement Project, MPKV, Rahuri for providing all facilities and research grants to carry out this experimental work and authors cited in references for providing necessary literature material.

## Conflict of Interest

The authors declare there are no conflict of interest relevant to this article.

## References

1. Bhalerao PD, Gaikwad GS. Productivity and profitability of *Bt.* cotton (*Gossypium hirsutum* L.) under various land geometry and fertilizer levels. *Indian J Agron.* 2010;55(1):60-63.
2. Chetana D, Mevada KD, Patel MV. Effect of low-cost management practices on yield, quality and economics of *Bt.* cotton (*Gossypium hirsutum* L.) under middle Gujarat conditions. *The Pharma Innovation J.* 2023;12(9):1495-1502.
3. Dadgale PR, Chavan DA, Gudade BA, Jadhav SG, Deshmukh VA, Pal S. Productivity and quality of *Bt.* cotton (*Gossypium hirsutum*) as influenced by planting geometry and nitrogen levels under irrigated and rainfed conditions. *Indian J Agric Sci.* 2012;84(9):1069-1072.
4. Dhillon RN, Chabra KL, Punia SS. Effect of crop geometry and integrated nutrient management on fiber quality and nutrient uptake by cotton crop. *J Cotton Res Dev.* 2006;20(2):221-223.
5. Elamathy S, Karunakaran KR, Saravanakumar V, Chandrakumar M. Performance of Indian cotton sector-an economic analysis. *Asian J Agric Ext Econ Sociol.* 2022;40(10):106-113.
6. Gohil VB, Patel DM, Chaudhary MM, Gohil NB, Patel JR. Yield maximization of *Bt.* cotton through agro-techniques. *Int J Plant Soil Sci.* 2023;35(23):650-657.
7. Kakade SU, Deshmukh V, Gawate AN, Rakhonde OS, Potdukhe NR. Effect of fertigation levels and canopy management practices on growth, yield and economics of *Bt.* cotton (*Gossypium hirsutum* L.). *The Pharma Innovation J.* 2023;12(6):4569-4575.
8. Kaul A, Deol JS, Brar AS. Response of different *Bt.* hybrids (*Gossypium hirsutum* L.) to canopy modification practices. *J Appl Nat Sci.* 2016;8:1188-1197.
9. Khetre OS, Shinde VS, Asewar BV, Mirza AB. Response of growth and yield of *Bt.* cotton to planting densities as influenced by growth regulators. *Int J Chem Stud.* 2018;6(4):485-488.
10. Munir MK, Tahir M, Saleem MF, Yaseen M. Growth, yield and earliness response of cotton to row spacing and nitrogen management. *J Anim Plant Sci.* 2015;25(3):729-738.

11. Parihar LB, Rathod TH, Paslawar AN, Kahate NS. Effect of high density planting system (HDPS) and genotypes on growth parameters and yield contributing traits in upland cotton. *Int J Curr Microbiol App Sci*. 2018;7(12):2291-2297.
12. Pendharkar AB, Solunke SS, Sawargaonkar GL, Kote GM. Response of Bt. cotton hybrids to different plant spacing under rainfed condition. *Adv Res J Crop Improv*. 2010;1(2):180-182.
13. Shekar K, Ramana MV, Ratna SK. Response of hybrid cotton to chloro-mepiquat chloride and detopping under high density planting. *J Cotton Res Dev*. 2015;29(1):84-86.
14. Shwetha NS, Halepyati AS, Pujari BT. Effect of detopping, removal of monopodia and plant spacings on nutrient uptake, quality parameters and economics of Bt. cotton (*Gossypium hirsutum* L.). *Karnataka J Agric Sci*. 2009;22(4):892-893.
15. Singh K, Singh HP, Rathore P, Singh K, Mishra SK. Manipulations of source sink relationships through mepiquat chloride for enhancing cotton productivity and monetary returns in north western India. *Indian J Cotton Res Dev*. 2017;31(1):62-68.
16. Uma Maheswari M, Murali Krishnasamy M, Kumar S, Sakthival N. Impact of high-density planting system and growth retardants on root growth and yield attributes in machine sown cotton. *The Pharma Innovation J*. 2019;80(3):123-130.