



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2022; 5(1): 13-18

Received: 06-11-2021

Accepted: 12-12-2021

**Parveen Kumar Jain**

School of Agriculture, Galgotias University, Plot No. 2, Yamuna Expressway, Opposite Buddha International Circuit, Sector 17A, Greater Noida, Uttar Pradesh, India

**Ganesh Datt Bhatt**

School of Agriculture, Galgotias University, Plot No. 2, Yamuna Expressway, Opposite Buddha International Circuit, Sector 17A, Greater Noida, Uttar Pradesh, India

**Mahesh Singh**

School of Agriculture, Galgotias University, Plot No. 2, Yamuna Expressway, Opposite Buddha International Circuit, Sector 17A, Greater Noida, Uttar Pradesh, India

**Corresponding Author:**

**Parveen Kumar Jain**

School of Agriculture, Galgotias University, Plot No. 2, Yamuna Expressway, Opposite Buddha International Circuit, Sector 17A, Greater Noida, Uttar Pradesh, India

## A study on saving water using drip irrigation technologies in paddy and enhancing yield

**Parveen Kumar Jain, Ganesh Datt Bhatt and Mahesh Singh**

**DOI:** <https://doi.org/10.33545/2618060X.2022.v5.i1a.90>

### Abstract

We know that water is essential for growing rice and it requires standing-water for proper growth and successful cultivation. The practice probably started when under prevailing conditions the farmers has tweaked rice seedling to grow in accumulated water. The first records of rice cultivation projects it as a dry seeded crop grown in ends of high land and forests cleared for shifting cultivation. It's in China that the process of puddling soil and transplanting seedlings began in low-lying area and river valleys. As transplanting gives a head-start for rice seedlings in their fight with weeds, this method took an easy route of acceptance in other parts of the world especially in South Asia where originally rice is grown under dry land conditions. But the current scenario is entirely looking for water saving technologies and the aim has been shifted from 'crop productivity' to 'irrigation/water efficiency'. This can be by micro-irrigation that can improve the water use efficiency," stated the survey. The issue of the growth of water-saving /intensive crops in the agriculture of the country. "Around >89% of groundwater pumped is used for irrigation. The paddy and sugarcane consume > 60% of irrigation water," said the survey. According to a report, as compared to the ratio of total used volume of water to the production quantity of world average: India's rice production is 2020:1325 m<sup>3</sup> per tonne (UNESCO 2010).

Paddy is a great guzzler, and as compared to flood prone plains of Indo-Gangetic plains of west Bengal and Bihar, paddy takes three times more water in Punjab due to availability of ground water and canal systems. Out of 5.0 million hectares area under cultivation in Punjab the major share is of rice (paddy and basmati 3.0 million hectares, whereas in contrast to 1.23 lakh ha of maize and 2.83lakh ha of cotton, Alike other crops paddy has been reported to be a successful drip feeded crop. Thus, rice cultivation is water intensive; it uses 1600 mm per season as per literature, (65 lakh l /ac). But in actuality much more volumes are applied in many delta areas in India (up to 2200 mm, 90 lakh l/ac).

The use of micro irrigation accompanied with fertigation has a benefit for rice production. At one side the water consumption is reduced to a great extent when we use drip and at the same time under non-stress situation paddy respond with higher yields. Irrespective of the planting technique, varieties, method, qclimate and location. While rice yields enhancement due to drip-fertigation ranged from 14.7% to 29.9%. (Over that in flood method) averaged upto 11.61 t/ha (2) ability in soil is kept close to field capacity and crop is grown to result in enhanced yields. It's possible by micro irrigation technologies.

The topic is reviewed in view to see the reported and works carried out by drip companies. We believe that results from such effort would lead to an actual extension possibility of that technology in the coming years. Of the 43.8 million ha paddy cultivated in India, even a 10% of it gets into drip irrigation that is 4.4 million ha. The impact would be multifold: Substantial saving of irrigation water; meaning irrigating more area under paddy with the available water. Substantial saving of energy; a boon power starved. Increased production of rice; a much-needed requirement for the country.

The paper will help to make an awareness that the paddy is not a waterlogged crop and can be successfully grown with Drip irrigation.

**Keywords:** Paddy, drip irrigation, micro irrigation, transplanted rice, yield

### Introduction

Currently, in India total food grain production has been estimated at record 291.95 mt which is 6.74 million tonnes higher production of food grain of total 285.21 mt achieved during 2018-19. (Abdelraouf, 2013) [2]. The rice production is at record estimated to 102.36 mt in kharif season (2020-21) crop year on the back of good monsoon rains and acreage, according to government data (FAO, 2010) [18]. The demand augmenting in parallel to growing population. India should be producing 1.7 mt of additional rice every year to ensure national food security (Das and Chaandra, 2013) [7].

By 2050, Demand for rice is expected to rise 25% it will have to produce 494 mt of food grains. That is unlikely to happen unless farmers adopt GAP-good agriculture practice, better farming techniques and utilize the inputs like fertilizers and water judiciously. Rice is an obvious focus for water saving as, it is grown on more than 50% of irrigation water and 30% of irrigated land (Barker *et al.*, 1999) [4].

"Irrigation capable to improves yields by 60-100% and is a real option to enhance food grain figures," says (Soman *et al.* 2018) [17]. The water saved by drip irrigating paddy - as much as 40 per cent of current usage - can be used to irrigate a larger area. This method of cultivation of rice utilizes more than 30 to 45% of the world's fresh water resources (Humphreys *et al.*, 2010) [12]. Normally in one hectare area under traditional practice 2000mm water is used on higher side as reported 1300-1600 mm [7].

"We want to increase the production without increasing water use or lowering quality," said Michele Conte, whose family has managed La Fagiana for decades and who has adopted the Netafim system on some of his land. In Asia, 17 mha of irrigated rice will have physical water scarcity and economic water scarcity respectively by 2025 (Tuong and Bouman, 2001) [6].

Rice is cultivated usually in a puddled condition with large volumes of water and grown in standing water. The water productivity is hardly 0.15 kg/m<sup>3</sup> a water, which is very low. Even in SRI (Sustainable rice Initiative) method the total water use is 745-800 mm per ha with standing water during the major crop phase. Aerobic rice is crops grown in non-puddled, well-drained, & non saturated soils without accumulating water (Bouman *et al.*, 2007).

A comprehensive package was tested that could reduce resource use and increase productivity in direct seeded rice which form 28% of the Indian rice cultivation. Jain Irrigation, the most highly water conscious Corporate in the country is contributing in its own way to food security by doing research and innovations in water saving and conservation technologies. The company has now tested and released an innovative method for cultivating rice on drip irrigated.

### Distribution of Paddy/Rice

The paddy second most important, food grain crop of the world, cultivated in as many as > 118 countries > 153.80 mha with a production of about 618.50 mt. The Asian region consumes >90% rice produced. The highest averaged yield productivity is reported from Egypt to the tune of 94 q/ha as against world's average of 40.4 q/ha. Table1 depicts the area, production and yield of major rice growing countries.

**Table 1:** Rice, acreage, production and productivity in major countries

Continent/Country	Area (m ha)	Production (m t)	Yield (t/ha)
Egypt*	0.65	6.35	9.84
USA*	1.35	10.47	7.78
Japan*	1.7	10.91	6.42
China	29.3	185.5	6.33
Vietnam	7.34	36.3	4.95
Indonesia	11.8	54	4.58
Myanmar	6.27	24.5	3.91
Asia	136.56	532.98	3.9
Bangladesh	11	40.1	3.64
Brazil*	3.73	13.28	3.62
Philippines	4.12	14.8	3.6
India	43	129	3
Pakistan	2.4	6.9	2.88
Thailand	10.2	27	2.65

Production values are for brown (unhusked) rice

**Source:** FAO Quarterly Bulletin of Statistics, 2005, Vol. 12(3/3):18-19 and \* FAO Production Year Book, 2004.

Approx. 41% of the annual food grain production area in India, about 12 million hectares are utilized for the paddy -wheat (Crop rotation) production system, It is grown in almost all the states, with Uttar Pradesh, West Bengal and Punjab leading in area, production and productivity, respectively. Table 2 depicts major states production data with respect to Paddy.

**Table 2:** Area, production and productivity of rice in various states of India (2020-21)

State	Area - Million Hectares	Production - Million Tonnes	Yield - Kg./Hectare
Andhra Pradesh	2.11	7.45	3540
Assam	2.47	4.73	1916
Bihar	3.34	8.24	2467
Chattisgarh	3.83	8.05	2101
Odisha	3.85	8.33	2160
Punjab	2.9	11.59	3998
Tamil Nadu	1.44	2.37	1642
Telangana	1.68	5.17	3075
Uttar Pradesh	5.99	13.75	2295
West Bengal	5.5	15.3	2784

**Source:** Maps of India. Com-2021

### Origin and History

Rice unlike other crops that man grows is intimately involved in the culture of numerous world societies the rice grain forms an integral part of rituals in Hindu and Buddhist religions. Hence the interlinking of rice with humans is not just as a commodity but as a substance that goes deep into the cultural milieu of the human race.

India and Burma/Myanmar are reported to be the center of origin of cultivated rice Vavilov (1926). Rice has been the first domesticated crop in Asia, as proved by the Sanskrit literature. Carbonized grains of rice were found in the excavations at Hastinapur (Uttar Pradesh) which belongs to 1000-750 BC. suggested. Chang (1985) reported the foothills of Himalayas and the Gangetic belt touching China. 24 various species are considered in *Oryza genus*.

### Precision farming of paddy

Precision farming is considered as modern technology of farming where 1) Timeliness of operations and 2) Precision in quantities of inputs and control measures are practiced. The different steps followed in precision farming varies from crop to crop and differences of these practices from conventional practices also varies from crop to crop.

In Precision farming of rice we recommend the following steps: Planting on raised beds, adopting a plant spacing of 0.2 m x 0.15 m, irrigating with drip following an irrigation schedule and fertilizer through fertigation scheduling and weed control by weedicide and chemigation for pest control.

### Cultivation Method

Based on experiments and field trials over a number of years, the cultivation method and drip system for rice should have the following components:

Adopt dry seeded aerobic method of rice cultivation Disc plough the field twice; first East-West and second time North-South directions For rainy season crop, it is advisable to prepare broad bed and furrow (BBF) system that will take care of drainage during heavy rain. For rabi and summer crops even flat seed bed is appropriate For BBF system for seeding; Bed width of 1 m, and height of 0.15 m and furrow width of 0.2-0.3 m Sow rice in rows on a broad bed or on flat seed bed. Rotavate to break clods

on the bed surface to achieve fine Mulch the seeded surface with rice husk.

### Rice varieties

Identification of varieties suitable for drip irrigation would take a large number of time-consuming experimental trials. However, it is an intelligent assumption that all the varieties so far found suitable for dry seeded cultivation would also be suitable for drip. Varieties tested under drip irrigation and fertigation by Jain irrigation.

### Varieties tested under drip fertigation

US 311, Arise 6129, SBH 999 (basmati), 25P25, 25P31, Try-R (2), BPT, Pusa sugandha (basmati), ADT -45, Pusa-2, WGL 32100.

### Types of rice cultivation

Paddy is mainly cultivated as uplands and lowlands, upland rice cultivations done with drill or broadcasting or can be cultivated by drip irrigated plots in line sowing. A good area under paddy in, Uttar Pradesh, Bihar, Orissa, Andhra Pradesh, Tamil Nadu is under rainfed /upland cultivation method. (Abhishek, 2020) [1].

The seed beds are maintained at adequate moisture with periodic drip irrigation in place of flooding. Another practice of Lowland rice cultivation with puddled soils, transplanting, or dapog/matt nursery method is done.

### Drip Irrigation - The concept

Drip irrigation is the slow, even application of water at low pressure to the root zone using a network of plastic tubing placed above the rooting zone (surface drip) or buried among the root branches inside the rhizospheres soil at a certain depth from surface. (Subsurface drip).

In drip irrigation method, crops are irrigated daily to the precise volume of water equivalent to the evapo-transpiration (ET) of the crop. It is estimated from daily Evaporation data using crop and canopy coefficients, the latter two factors vary with the age of the crop and the size of its canopy. These are the two factors that affect the volume of transpiration of the plant/ crop that changes with growth of the crop. Factoring in of these two coefficients is what makes the water requirement estimate unique to that particular crop at that particular stage of its growth.

### Sowing and germination Irrigation

It is essential to pre irrigate and fully wet the broad bed before sowing the seeds. Keep the bed uniformly wet till germination and crop establishment.

### Mulching

One of the main issues in dry seeded and drip irrigated rice is the germination and growth of weeds. The standing water in the conventional flooded rice will suppress the weed germination. We recommend use of rice husk mulch on the beds after seeding. Mulch is applied 2-3 cm thick on the bed surface. Besides reducing weed growth it also helps reduce evaporation from the soil surface. However, this is an optional recommendation.

### Weedicide application

Use of weedicide to prevent weed growth is also plausible. Studies have shown that a pre-emergence weedicide, Pretilachlor sprayed to the seed bed at 1250 ml/ha rate within 72 hours of sowing controls weed infestation effectively.

### Fertilizer application through Fertigation

Use of drip technology provides a golden opportunity to apply nutrients as per the need of the crop at each growth stages. In rice also this technology enhances nutrient use efficiency.

Water is not the only need of the plant. To uptake this water efficiently, it requires proper air-water balance within the root zone. Drip irrigation, with its low application rate, prevents the saturation of water within the root zone and continuously maintains field capacity. This provides a favorable condition for the growth of the plant. Drip irrigation also helps to use fertilizer efficiently. With drip irrigation water can be provided at frequent intervals which helps maintain required soil moisture level within the vicinity of the plant roots.

### Crop Rotation

Drip irrigated rice can be followed by drip irrigated wheat. The rotation crops can be, a second rice, mustard, vegetables, or seed spices. Both the rotation crops are adjusted in their spacing to suit the drip system on the ground.

### Fertigation schedule

**Table 3:** Fertigation schedule adopted in drip irrigated rice in AP\* (Fertilizer dose 180:80:80 kg NPK per acre) \$DAP-days after Germination \* Schedule will vary with location.

	Urea	SSP	MOP	ZnSO <sub>4</sub>	Schedule (urea) kg/day/ac	Schedule (MOP) kg/day/ac
Basal (soil)		500	0	10	-	-
till 10 DAPS	65	0	0	-	6.5	-
11-35 DAP	196	0	16	-	7.8	0.64
36-55 DAP	65	0	17	-	3.3	0.85
55-65 DAP	65	0	50	-	6.5	5
65- 70 DAP	0	0	17	-	-	3.4

### Integrated Nutrient management


- Incorporate green manures/bio fertilizers
- Nitrogen management by leaf color chart (LCC)
- Practice fertigation daily.

The different fertilizers are applied in foliar form with water solubilizing fertilizers in periodic intervals under fertigation, but basal is applied in the seed beds while making the beds.

### Water Management

By reducing infiltration, seepage, percolation, and evaporation, efficient water use during field preparation in upland aerobic rice is lowered to the tune of about 51% and 32-88% higher water use efficiency, than flooded rice (Bouman *et al.*, 2005) [5]. The labor use is also saved in aerobic rice as more labor is required for puddling, land preparation, transplanting, and irrigation activities in flooded rice (Wang *et al.*, 2002) [19].






**Drip Irrigation - The concept**

The concept of using each drop of water in a useful way and that leads to production is the aim behind Drip irrigation. Earlier it was considered that only few crops can be grown with drip and with time many new crops brought under drip but crops like

paddy are still be considered as non-drip crop, the crop is responsible for heavy water use and also responsible for 10% methane leading to Green house effect. The earlier experiments by drip companies like Jain Irrigation and Netafem resulted in at par production of rice. (GOG, 1994) [10].



**Fig 1:** Drip irrigated paddy layout

Drip irrigation means the slow, even application of irrigation water at low pressure to the root –zone using a network of plastic tubing placed above the rooting zone (surface drip) or buried among the root branches inside the rhizospheres soil at a certain depth from surface. (Subsurface drip).

In drip irrigation method, crops are irrigated daily to the precise volume of water equivalent to the evapo-transpiration (ET) of the crop. It is estimated from daily Evaporation data using crop coefficients and canopy coefficients, the latter two factors vary with the age of the crop and the size of its canopy these are the two factors that affect the volume of transpiration of the plant/crop that changes with growth of the crop. Factoring in of these two coefficients is what makes the water requirement estimate unique to that particular crop at that particular stage of its growth (Soman *et al.* 2018) [17].

Govindouram named Parthasarathy and 23 other farmers has used Drip on Paddy, onion and Maize. He observed a 15-20 percent increase in yield and also water saving 30-40 percent. He got Innovative Farmer Award for this in 2015. (Rao, 2003). 20% of global white rice area is in India make it to cover largest area and output 42% of total grains produced, the flood irrigation is practiced in 85% area, so if this 50% area out of this be brought in the Drip will be a great contribution in water saving. (Ahmad *et al.* 2004) [3]

**Sowing and germination Irrigation**



**Fig 2:** Paddy grown with drip irrigation

It is essential to pre irrigate and fully wet the broad bed before sowing the seeds. Keep the bed uniformly wet till germination and crop establishment. And after that a periodic irrigation is

**Table 4:** Water requirement in rabi paddy

Month	Pan E (mm/day)	Total water Reqt. (lit/Day/ha)
Feb 16-28	6.68	18704
Mar 1-15	6.73	40044
16-31	6.82	47740
April 1-15	6.09	56840
16-30	6.53	60947
May 1-15	7.44	70019
16-31	7.57	58878

A farmer from the Tirupur district of Tamilnadu of village

prescribed as water IR scheduling.

Some advance work under funded projects from Israel are also going in TNAU titled, 'Feasibility of growing rice under drip irrigation', the experiment has been taken close to the 'kuruvai' season and scientists have planted four varieties 'TRY II', 'ADT 45', 'ADT 36' and 'Anna (R) 4' for ascertaining the growth of paddy in saline fields on the campus of the institute (Puranik *et al.*, 1992) [14].

The institute, located in the Manikandam block, has taken up various research programmes on evolving paddy varieties suitable for sodic soil. Netafim, an Israeli company specialising in drip irrigation technique, had sanctioned Rs. 4 lakh to the institute to take up the research, said P. Pandiyarajan, Dean of the Institute.

Speaking to The Hindu after inaugurating the programme, Mr. Pandiyarajan said that this was the first time the institute had taken up the research using drip irrigation technique, although similar experiments had been conducted at Aduthurai and other places. About 2 kg of seeds was used for each variety. He said that the experiment would include study on nutrient management, irrigation management and good cultivation practices. The entire 60 cents of the field has been covered with over 77 laterals, each of 50 metre in length. "Each lateral contains 130 holes or drip spacing through which water and water-soluble fertiliser will be supplied," he said (Rangachari *et al.*, 2000) [15].

### Sustainable Rice Production

With drip- fertigation one achieves a sustainable method for rice production. Conservation of water, energy and fertilizer while enhancing yield by 1 to 1.5 ton per ha or more results in a Win-Win situation. We produced 22% more rice using 66% less water; (water use in drip is 8 million liter per ha compared to flood irrigation where it is 23.75 million liter per ha) and using 52% less power for pumping water. More rice is produced per unit input; Water productivity also improved to 500 to 1000 g/1000 liter from a figure of 120 to 290 g/1000 liter.

**Table 5:** Sustainable use of water resources

Irrigation	Flood	Drip
Area, ha	1	1
Water Use, m <sup>3</sup> /ha	23,750	8,000
Yield, kg/ha	7,000	8,500
WUE (Water Use Efficiency) kg/m <sup>3</sup>	0.29	1.06
Selling Price of Rice/Kg	20	20
Income/acre	1,40,000	1,70,000
Value Creation Efficiency/m <sup>3</sup>	5.89	21.25

S Improving yields of rice is imperative to addressing growing food security challenges S Improves water use efficiency of rice crop and the value of water. The value creation efficiency in flood method is only 5.89 per 1000 liter water compared to 21.25 per 1000 liter in drip.

S Smallholder farmers have an impact on biodiversity and ecosystems and are dependent on them S At the same time, improving their practices and water productivity can help better preserve and protect natural resources S Improves yield, income and reduces water, energy and fertilizer use of the follower crop after rice also. Thus drip-fertigation ensures water, energy and food security in a sustainable manner.

### Harvesting paddy

Various Paddy varieties normally mature in 95-160 days. The visual symptoms that the crop is ready for harvesting proper

stage for harvesting is yellow coloring and ripening of about >80% panicles of the area. The harvest is of good quality specially when harvested at proper stage and time in a dry weather. These days harvesting with combine harvesters is quite common and fast and time saving.

### Threshing operation

The thrashing practice with manual operation is still common in country where the rice threshing is done in the farm itself. But now a days combine threshers are available which do all three operations simultaneously and that too time saving.

The produce moisture content for storage and for selling in mandis of rice grains is 12%.

### Conclusion

Drip Irrigation or micro irrigation has been proved as one of the best water conservation technology for heavy consumption of water in rice growing, that too without scarifying the yield. The aerobic rice can be successfully grown just like all other crops with drip by maintaining moisture at adequate level in root zones

The investment cost of drip irrigation is a concern that cost between Rs 30,000 - 70,000 per acre. The initial cost of installation is beyond afford capacity of marginal farmers (86%) and cannot afford this. In many states under centre or state Govt schemes subsidies to the tune of 60-90% against beneficiary share of 10-40%. In many cases this beneficiary share is also adjusted in digging trenches or civil works by the service provider companies.

There is a need to provide subsidies to farmers so that the MIS systems can be promoted in rice system also.

### References

1. Abhishek. Rice Cultivation Guide: From Introduction to Yield. Agriculture review. 2020, 2.
2. Abdelraouf RE, Habbasha S, Taha MH, Refaie KM. Effect of irrigation water requirements and fertigation levels on growth, yield and water use efficiency in wheat. Middle-East J Scientific Res. 2013;16:441-450.
3. Ahmad MD, Masih I, Turrall H. Diagnostic analysis of spatial and temporal variations in crop water productivity: A field scale analysis of the rice-wheat cropping system of Punjab, Pakistan, Journal of Applied Irrigation Science, 2004;39(10).
4. Barker R, Dawe D, Tuong TP, Bhuiyan SI, Guerra LC. The outlook for water resources in the year 2020: challenges for research on water management in rice production. In: Assessment and Orientation Towards the 21st Century, Proceedings of 19th Session of the International Rice Commission, Cairo, Egypt, 7-9 September 1998. FAO, Rome. 1999, 96-109.
5. Bouman BAM, Peng S, Castaneda AR, Visperas RM, Yield and water use of irrigated tropical aerobic rice systems. Agricultural Water Management. 2005;74:87-105.
6. Bouman BAM, Tuong TP. Field water management to save water and increase its productivity in irrigated rice. Agricultural Water Management 2001;49(1):11-30.
7. Dass A, Chandra S. Irrigation, spacing and cultivar effects on netphotosynthetic rate, dry matter partitioning, and productivity of rice under system of rice intensification in mollisols of northern India. - Expl. Agri. 2013;49(4):504-523.
8. FAO (Food and Agriculture Organization of the United Nations), Electronic on-line database. 2010. Online at

<http://www.fao.org>.

9. Ghosh BN, Singh RD. Effect of conjoint use of farmyard manure and rice (Oryza sativa) –wheat (Triticum aestivum) system in Uttaranchal mid-hill soils. *India J. of Agric. Sci.* 2003;73(12):680-683.
10. GOG. Drip irrigation in India, Indian National Committee on Irrigation and Drainage, Ministry of Water Resources, Government of India, New Delhi. 1994.
11. Hegde DM, Srinivas K. Growth, yield, nutrient uptake and water use of banana crops under drip and basin irrigation with N and K fertilization, *Tropical Agriculture (Trinidad)*, 1989;68:331-334.
12. Humphreys E, Kukal SS, Christen EW, Hira GS, Singh B, Yadav S *et al.* Halting the groundwater decline in north-west India—Which crop technologies will be winners? *Adv. Agron.* 2010;109:155-217. doi:10.1016/B978-0-12-385040-9.00005-0.
13. Janpen P, Minamisawa K, Teamtai song K, Boonkerdand N, Teamroong N. The communities of endophytic diazotrophic bacteria in cultivated rice (Oryza sativa L). *Applied Soil Ecology.* 2009;42:141-149.
14. Puranik RP, Khonde SR, Ganorkar PL. Constraints and problems as perceived by non-adopters and adopters of drip irrigation system. *Agricultural Situation in India.* 1992; XLVII(1):33-34.
15. Rangachari R, Nirmal Sengupta, Ramaswamy R Iyer, Pranab Banerjee, Shekhar Singh. Large Dams: India's Experience, Final Report, prepared for the World Commission on Dams, Cape Town, South Africa, Secretariat of World Commission on Dams. 2000.
16. Rao PS. Study on efficient use of irrigation water device: a case study. *Indian Journal of Agricultural Economics*, July-September. 2003;58(3):504.
17. Soman P *et al.*, Effect of Drip Irrigation and Fertigation on the Performance of Several Rice Cultivars in Different Rice Ecosystems in India. *International Journal of Agriculture Sciences*, ISSN: 0975-3710 & E-ISSN: 0975-9107. 2018;10(14):6672-6675.
18. Tuong TP, Bouman BAM. Rice production in water-scarce environments, In: *Proc. Water Productivity Workshop*, 12-14 November 2001, Colombo, Sri Lanka. International Water Management Institute, Colombo, Sri Lanka. *International journal of Agronomy and Plant Production.* 2003;3(7):241-254.
19. Wang HQ, Bouman BAM, Zhao DL, Wang CG, Moya PF. Aerobic rice in northern China: opportunities and challenges. In B. A. M. Bouman, *et al.*, eds. *Water-wise-rice production. Proceedings of the international workshop on water-wise rice production*, 8-11 April 2002, Los Banos, Philippines. International Rice Research Institute, Los Banos. 2002, 143-154.
20. Yadav BD, Joon RK, Kumar Virender. Response of early maturing guar variety to date of sowing and seed rate. *Adv. Arid Legumes Res.* 2001, 199-202.