



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2022; 5(1): 26-31

Received: 10-11-2021

Accepted: 16-12-2021

**Zigale Semahegn**

Ethiopian Institute of Agricultural  
Research, Melkassa Agricultural  
Research Center, National  
Sorghum Research Program,  
Ethiopia

## Intercropping of cereal with legume Crops

**Zigale Semahegn**

### Abstract

The technique of farming two or more crops in the same field at the same time is known as intercropping. It has been practiced for decades. It provides a number of benefits, including more efficient and effective resource utilization, less soil erosion, increased yield output on a limited cultivated field, risk reduction for smallholder farmers, and higher income per unit of work during times of labor scarcity. Intercropping legumes with cereals, in general, allows for more energy-efficient and continuous agriculture. However, as the world's population grows, a lack of arable land is a major developmental limitation in many rising countries' capacity to meet their basic food and nutritional demands. The most popular types of intercropping used to overcome poorly farmed terrain are mixed intercropping, row intercropping, and strip intercropping. Intercropping is influenced by a variety of factors, including the choice of compatible species, the timing of establishment/planting, knowledge of the physiology of the species to be cultivated at the same time, their growth habits, plant population density, canopy and root architecture, and water and nutrient usage. Various competition indices in cereal-legume intercropping have been used to measure crop yield and efficiency per unit area of land. Only a few of these are the Land Equivalent Ratio, Area Time Equivalent Ratio, Aggressivity, Relative Crowding Coefficient, Competitive Ratio, and Actual Yield Loss. A well-managed cereal-legume intercropping system has been found to have beneficial impacts on resource utilization and combined production of low-input crops in the past. The most useful strategy for sustainable agriculture and food security is to address the cereal-legume intercropping system to all farmers in developing countries.

**Keywords:** Intercropping, cereal-legume intercropping, benefit, effect

### Introduction

Intercropping is a low-input farming strategy that involves cultivating two or more crops in the same farmed area at the same time to improve resource efficiency and cropping system flexibility. It's also an old and popular planting approach that tries to match crop preferences with growing resources and manpower available. Smallholder farmers in Asia, Africa, and Latin America have long used intercropping, and it is gaining popularity due to its ability to produce high yields with minimum inputs and its ability to save space (Yu *et al.* 2015; Martin-Guay *et al.* 2018; Li *et al.* 2020a, b) <sup>[50, 25, 23]</sup>. Pigeonpea has been shown to have the greatest potential for intercropping with cereals in Eastern and South Africa due to its drought-tolerance features, and it plays a significant role in family nutrition, output, and revenue. Pigeonpea thrives in hard, drought-prone circumstances, unlike most legumes, and is commonly intercropped with sorghum, maize, millet, and other cereal crops. The Konso people, who live in an arid pastoral area of south-central Ethiopia, have intercropped traditional pigeonpea with ratooned sorghum (Westphal, E. 1974) <sup>[45]</sup>. Monocropping has a number of disadvantages such as increasing artificial fertilizer use, soil fertility loss, increases the spread of diseases and insects. when compared to intercropping. It increases output across a limited area of cultivated land, minimizes soil erosion due to fast soil cover in intercropping conditions, lowers risk for smallholder farmers, and provides more money per unit of effort in labor shortages (Anil L, Phipps RH, 1998.). Intercropping legumes and grains is a popular technique in many parts of the world, and it can serve as a model for farmers with limited resources who want to maintain output and food security. Consider the following aspects while intercropping: compatible species, planting timing, understanding the physiology of the species to be intercropped with, their growth habits, canopy and root architecture, and water and fertilizer consumption.

Intercropping has a higher biotic efficiency than monocropping due to the larger soil mass explored. In previous decades, this advanced agricultural approach was successful in achieving agriculture's goal (Ofori, 1987) <sup>[30]</sup>.

**Corresponding Author:**

**Zigale Semahegn**

Ethiopian Institute of Agricultural  
Research, Melkassa Agricultural  
Research Center, National  
Sorghum Research Program,  
Ethiopia

Several studies on cereal-legume intercropping systems have been undertaken (Dereje *et al.*, 2016; Tenaw *et al.*, 2016; Hailu, 2015; Gebremichael *et al.*, 2019; Tamiru *et al.*, 2013; Alemayehu *et al.*, 2018; Addisu *et al.*, 2014; TANA & MULATU, 2000; Dahmardeh, *et al.*, 2010)<sup>[12, 40, 19, 17, 38, 5, 2, 39, 13]</sup>. High population growth and a scarcity of cultivated land are major developmental issues in many emerging countries, making it difficult to meet their food and nutritional needs. Weeds, drought, and nutrient mining are also major factors in low agricultural output in underdeveloped nations, in addition to inadequate soil fertility and limited access to resources for small-scale farmers (McCann, 2005). Intercropping is more productive than Ethiopia's traditional cereal-based cropping methods (Tesfa Bogale *et al.*, 2002). As a result, the intercropping system is critical for increasing crop production, enhancing soil fertility, reducing weed computation, increasing profit, and lowering crop failure risk. Farmers, on the other hand, cultivate their land without using soil and water conservation measures for lengthy periods of time, resulting in extremely deteriorated soil. Farmers are also hesitant to use man-made fertilizers due to their expensive cost and lack of information, resulting in low crop yields in a given amount of land. To address these issues, it is critical to do and disseminate research on soil and water conservation, as well as fertilizer application, particularly to increase productivity. They also use an intercropping strategy to help sustain agriculture by improving soil fertility in the country. Intercropping is an important part of subsistence farmer crop production methods, hence a review on cereal-legume intercropping was initiated.

### History of intercropping

Intercropping is a long-standing tradition, probably dating back to the beginning of civilized agriculture. However, there is no precise date for when the first intercropped field appeared, but historians (Anders *et al.*, 1996)<sup>[7]</sup> claim that intercropping existed early in the genesis of agriculture. Furthermore, despite the difficulties in pinpointing the exact dates when intercropping in the form of mixed garden plots initially appeared, it has been proved that they were common during Paleolithic times (Plucknett and Smith 1986)<sup>[32]</sup>. The transition from informal to formal mixed garden plots did not happen overnight; it was a long process that began in Paleolithic times.

In many places of the world, intercropping methods have long been established (Francis 1986)<sup>[15]</sup>. Intercropping is crucial in the tropics, and it is still commonly used. In terms of crops and systems, there is a lot of natural genetic variation in these areas (Anders *et al.*, 1996)<sup>[7]</sup>. Farmers' use of intercropping falls as the temperature and rainfall drop (Harris 1976)<sup>[20]</sup>. This trend is attributable to a drop in the number of plant species that have been adapted to tough growing conditions, as well as farmers' preference for species that have a better chance of producing something even if the season is terrible.

Intercropping began to wane in many industrialized countries as a result of mechanization and specialization of new commercial and production commodities with the introduction of sophisticated agriculture. Individual commodities were studied as independent components of the agricultural system, with a focus on crop species and cultivars. The most effective way for raising agricultural output was thought to be specialization. With a single commodity, this could have worked.

### Main Types of Intercropping

The technique of planting two or more crops on the same piece of land at the same time is known as intercropping.

Intercropping uses temporal and spatial crop intensification to meet the needs of farmers and the growing environment (Eskandari *et al.*, 2009)<sup>[14]</sup>. Furthermore, while intercropping, the component species planted throughout the crop season compete for available resources. The following are three types of intercropping systems that are commonly accepted in different nations (Ofori *et al.*, 1987)<sup>[30]</sup>.

#### Mixed Intercropping

Mixed intercropping is when two or more crops are planted together without a set row design. On occasion, it's also referred to as mixed cropping (Von Cossel *et al.*, 2019)<sup>[44]</sup>. Grass-legume intercropping is the best example of mixed intercropping in a pasture-based cropping system (Gulwa *et al.*, 2017)<sup>[18]</sup>. In locations with limited land resources, mixed intercropping is primarily employed to meet food and forage demands (Undie *et al.*, 2012).



Source: Downloaded from Google

Fig 1: Mixed Intercropping

#### Row Intercropping

Row intercropping is a type of intercropping in which one or more crops are planted in rows and intercrops are cultivated simultaneously in a row or without a row. Row intercropping is a common technique for increasing resource efficiency and productivity (Varma, 2017)<sup>[42]</sup>.



Source: Downloaded from Google

Fig 2: Row Intercropping

#### Strip Intercropping

Strip intercropping is a type of intercropping in which two or more crops are planted simultaneously in strips on sloppy soil. It is typical to increase radiation efficiency in isolated and underprivileged places (Yang *et al.*, 2015). A range of soil saving and depleting crops are planted in alternate strips running perpendicular to the slope of the ground or the direction of prevailing winds. Strip cropping has two main goals: reducing soil erosion and boosting yield output from low-fertility soils.



Source: Downloaded from Google

Fig 3: Strip Intercropping



### Relay intercropping

Relay intercropping is the practice of growing two or more crops at the same time for a portion of each crop's cultivating time. The following crop is sown after the preceding crop has completed a significant portion of its life cycle and has reached reproductive stage or approaching maturity. When there is a paucity of time and soil moisture, relay cropping is the best option (Balde *et al.*, 2011) [87]. Before the previous crop is harvested, the second crop is planted, and both crops remain in the field for a period of time. Sequential cropping, on the other hand, yields fewer seeds than conventional planting, requiring more seeds of the successive crop to maintain a productive stand.



Source: Downloaded from Google

Fig 4: Relay Intercropping

### Major aspect to be considered in cereal-legume intercropping system

In order to develop an intercropping system, many elements must be considered both before and during the growing period (Seran and Brintha, 2010) [35]. Crop population density, light interception, crop variety, and nutrients, for example, all influence the potential of a cereal-legume intercropping system to produce nitrogen (Francis, 1989). Regardless, the plant's location, light, growth habit, water, and fertilizer use all influence compactable crop selection (Brintha and Seran, 2009) [35].

### Competition indices in intercropping

The following indicators were used to determine the degree of competition between different intercropped crops (Connolly *et al.* 2001; Weigelt & Jolliffe 2003). The Land Equivalent Ratio (LER) is the most well-known technique for comparing biological efficiency and yield per unit area of land in comparison to monocropping systems; when the LER is more than 1.0, it signifies that intercropping yielded more than growing alone (Onwueme and Sinha, 1991). Allen and Obura (1983) [6], for example, evaluated 1.22 and 1.10 LERs for maize-soybean intercropping over two years and concluded that intercropping outperformed monocropping. According to Abera *et al.*, the LER values varied from 1.15 to 1.42, indicating that maize-bean intercropping had higher productivity and land use efficiency in terms of crop output per unit area than solo planting (2005). In most situations, intercropping exceeded lone cropping, with Land Equivalent Ratios (LERs) often exceeding one (Osman *et al.*, 2011) [31]. The Area Time Equivalent Ratio (ATER) is a second competition indices approach that provides a more precise assessment of the yield benefit of intercropping with monocropping in terms of time consumed by component crops in intercropping systems (Hiebsch & McCollum 1987; Aasim *et al.* 2008; Yahuza 2011). It can be calculated using the formula  $(ATER) = LER \times D_c / D_t$  (Heibsch, 1980), where LER is the crop's land equivalent ratio,  $D_c$  is the crop's time, and  $D_t$  is the overall system's time. The third method is aggressivity (A), which is used to assess the competitive relationship between two crops in a mixed cropping system (Willey, 1979) [46]. The relative crowding coefficient (K), which is a measure of one species' relative dominance over another in a mixture, is the

fourth approach (Banik *et al.*, 2006) [10]. Because it is based on yield per plant, the actual yield loss (AYL) index gives more exact information on competition between and within component crops, as well as the performance of each species in the intercropping system, than the other methodologies (Banik *et al.*, 2000) [9].

### Intercropping system and its benefit

Intercropping is a practice of growing two or more crops on the same piece of land at the same time to promote agricultural sustainability. Among Ethiopia's poor farmers, it's a regular practice. Maize-beans, sorghum-cowpea, millet-groundnuts, maize-cowpea, maize-soybean, maize-pigeonpea, maize-groundnuts, and rice-pulses are all common intercropping crop combinations. The features of an intercropping system vary depending on the soil, crop species, climate, economic position, and preferences of local farmers. Intercropping systems, which largely intercrop cereal and legume crops that enhance soil fertility and sustainability, reduce pests, diseases, and weeds and promote soil-physical conditions by receiving more radiation and making better use of available water and nutrients (Willey *et al.*, 1983; Horwith, 1985; Ofori and Stern, 1987; Jarenyama *et al.*, 2000; Sanginga and Woome, 2009) [30, 33]. Intercropping, rather than monocultures, provides year-round ground cover, preventing soil desiccation and erosion. Farmers can boost water use efficiency, protect soil fertility, and prevent soil erosion by planting multiple crops on the same piece of land at the same time, which are the main problems with mono-cropping. Intercropping legumes and cereals improves legume nitrogen fixation while also providing residual nutrients to the soil (Sarkar *et al.*, 1995) [34]. The cropping system's effective use of critical resources is governed by the inherent efficiency of the individual crops that make up the system, as well as the complementary influence of the crops (Willey and Reddy, 1981).

### Effect of Intercropping on Cereal Crops

In semi-arid regions of the world, intercropping legumes and grains is common. In tropical agriculture, legume-cereal intercropping is the most well-known intercropping approach, and it can reduce the amount of nutrients absorbed from the soil as compared to monocropping (Synadon and Harris, 1979) [36]. Legumes will fix nitrogen from the environment in the absence of nitrogen fertilizer and will not compete with cereals for nitrogen fertilizer (Gyamfi *et al.*, 2007) [4]. When compared to monocropping, intercropping legumes and cereals boosts productivity and soil fertility. Smallholder farmers are increasingly intercropping grains and legumes due to the legume's ability to minimize soil erosion and reduce soil fertility.

### Effect of intercropping on light interception and radiation use efficiency

The most valuable factor for crop growth and development was light, which varied from other growth factors in that it is only available in the present and hence must be taken straight away, whereas other resources are frequently pools waiting to be used by plants (Willey, 1979) [46]. Plant population density and row organization are thought to affect light transmission through the canopy (Jaya *et al.*, 2001) [21]. They also discovered that planting maize with a N-S orientation at an average density of 7.1 plant m<sup>-2</sup> lowered canopy maximum temperatures by 1.2°C at 40 cm above ground. Reduced irradiance was linked to a 70 percent drop in temperature; the drop in temperature, in particular, was

highly sensitive to row orientation and plant density, and in some circumstances resulted in an increase in temperature. With an irradiance of above 300 Wm<sup>-2</sup> at mid-day approximately 5 weeks after planting, a plant density of 7 plants m<sup>-2</sup> at N-S orientation was found to be promising for cauliflower-maize intercropping in arid lowland situations (Jaya *et al.*, 2001) [21]. This must be synchronized with cauliflower development, as early growth of cauliflowers necessitates higher irradiances to maintain enough glucose supply.

Light interception and light use efficiency are terms that can help you understand how cropping systems, particularly intercrops, obtain and use resources. Increased production could result from increased sunlight/solar radiation interception, increased light use efficiency, or a combination of the two (Willey, 1990) [48]. When two crop species are mixed and cultivated together rather than separately, light interception improves on a regular basis, either as a result of a longer time of soil covered (temporal advantage) or a more comprehensive soil cover (spatial advantage) (Keating and Carberry, 1993) [22]. Because the total crop population density of intercrops is higher, they can capture more light, especially early in the growing season. Intercrops with non-synchronous canopy development patterns and variable maturation times can have more leaf area during the growing season and produce more fruit. Intercrops, according to Carandang (1980), increase light interception by 30-40%, allowing for more efficient use of sunlight.

The overall system's light interception is detected by the geometry of the crop and the layout of the vegetation (Trenbath, 1983). By requiring the earlier crops to be put between properly broader rows of the taller once, intercropping between large and tiny canopy crops optimizes light interception (Seran and Brintha, 2010) [35]. Two characteristics that influence yield in proportion to occurrence radiation in an intercropping system are the total amount of light intercepted and the efficiency with which intercepted light is converted to dry matter (Keating and Carberry 1993) [22]. The amount of radiation intercepted in maize-bean intercropping was larger than in a single crop, according to Tsubo, Walker, and Mukhala (2001).

#### **Effect of intercropping on nutrient use efficiency**

Despite the challenges of determining helpful or competitive effects, nutrient utilisation in intercropping systems has gotten a lot of attention (Morris and Garrity 1993a) [27]. Due to variances in nutrient uptake techniques among crop species and between individual nutrients, quantifying competitive implications is difficult. More nutrient use can be seen in intercropping systems, both spatially and temporally. Spatial nutrient uptake can be maximized by increasing root mass when crops in an intercropping system have extremely high nutrient needs at various periods, whereas temporal nutrient uptake can be found when crops in an intercropping system have extremely high nutrient demands at different times (Anders *et al.*, 1996) [7]. Cereal-legume intercropping systems exhibit distinct rooting and nutrient uptake forms than monocrops, indicating that more effective use of available nutrients and increased N-uptake in the intercrop may be found (Fujita and Ofosu-Budu, 1996) [16]. Because all roots have the same direction and depth below the surface when just one species is growing, they tend to compete with one another (Seran and Brintha, 2010) [35]. Certain research from outside of Sub-Saharan Africa have demonstrated the relative efficiency of intercrops over monocrops. Maize and cowpea intercropping is beneficial on nitrogen-deficient soils, according to Vesterager, Neilsen, and Høgh-Jensen (2008) [43], and maize-cowpea intercropping increases nitrogen, phosphorus,

and potassium content when compared to monocrops of maize, according to Dahmardeh, Ghanbari, Syahsan, and Ramrodi (2010) [13]. Intercropping may speed soil nutrient depletion, notably for phosphorus, as a result of more effective soil nutrient usage and higher removal through harvested crops, in addition to the favorable effects on cereal crops (Mucheru-Muna *et al.*, 2010) [28]. Chalka and Nepalia (2006) [11], on the other hand, discovered that intercropping maize with soybean reduced NPK depletion while increasing N uptake. In order to reduce the use of foreign inputs, recent projects in Africa to improve soil fertility have included the use of legumes as an intercrop and/or in rotation (Sanginga and Woomer, 2009) [33].

#### **Effect of intercropping on weed management**

Weeds compete for agricultural growth and development resources, reducing crop output quantity and quality (Memon *et al.*, 2013). Furthermore, weeds that promote the spread of disease and insect pests, both of which can reduce agricultural productivity. To reduce the impact of weeds, farmers have used a range of strategies. Two of the most frequent ways for suppressing or controlling weeds are herbicide spray and hand weeding. Herbicide treatment, on the other hand, has negative consequences for environmental resources such as soil, water, and beneficial insects, as well as increasing production costs and fostering herbicide resistance (Santos *et al.*, 2014). As a result, research has demonstrated the value of agronomic options for weed control, such as intercropping, in reducing the negative effects of weeds as well as the side effects of herbicide treatments (Shah *et al.*, 2011; Khan *et al.*, 2013). Conventional intercropping systems are one of the possibilities for weed suppression or control as compared to solo crops (Ananthi, 2017) [37]. In actuality, weed growth is reliant on all component crops' competitive abilities, which in intercropping is highly reliant on the competitive abilities of the crops community and their respective plant densities (Willey *et al.*, 1980) [47]. For example, because of the legume soil cover, which created an inappropriate environment, the most frequent type of weed, striga infestation, has been considerably reduced (Musambasi *et al.*, 2002; Mashingaidze, A.B., 2004) [39, 26]. According to Chalka, M.K. and V. Nepalia (2006) [11]. Bean-maize intercropping reduced weed biomass by 50-66 percent when sown at a density of 222,000 plants ha<sup>-1</sup> for beans, which is 33 percent of the maize density (37,000 plants ha<sup>-1</sup>).

#### **Residual effects of cereal-legume cropping system**

Decomposing roots and falling leaves give nitrogen and other nutrients for the following crop once the intercrop is harvested (Geno and Geno, 2001). The residual influence of the pulse crop on the next crop is highest when the remaining pulse crop is left on the field and tilled under after harvest. When a substantial amount of nitrogen is left in the grain harvest, however, the land loses more nitrogen than the pulse crop can replace (Geno and Geno, 2001). The intercropped legume may store nitrogen in the soil that will not be useful until after the planting season, so increasing soil fertility and the value of a subsequent crop (Ofori and Stern 1987; Ledgard and Giller, 1995) [30]. Maize grain output, for example, was 46 percent higher when soybeans were planted following maize and natural fallow (Yusuf *et al.*, 2009). According to Herridge (1990), maximizing the total amount of nitrogen in the legume crop, the proportion of nitrogen derived from N<sub>2</sub> fixation, the proportion of legume N mineralized, and the efficiency with which this mineral N is utilized is required to increase the contribution of legume nitrogen to a subsequent crop. Unfortunately, it is not always possible to optimize these values.

## Summary and Conclusion

Intercropping is a popular agricultural method that involves growing two or more crop species in the same cultivated area at the same time. Cereal-legume intercropping is an old and frequent practice in developing nations, and it might be one of the most important factors in ensuring sustainable agriculture and food security for resource-constrained farmers. In developing countries, major intercropping systems such as mixed, row, strip, and relay intercropping are used. The developing world's key hurdles in meeting their food and nutritional needs include high population growth and shortage of land resources. In addition, low agricultural output in developing countries is caused by declining soil fertility, limited access to resources for impoverished farmers, weeds, drought, and nutrient mining. Cereal-legume intercropping has a number of advantages, including improved resource efficiency, reduced soil erosion, increased soil fertility, weed and pest control, increased yield production on a limited cultivated land, risk reduction for poor farmers, and higher income per unit of labor during times of labor scarcity. The choice of suitable crop species, planting/establishment time, knowing the physiology of the crop species, their growth habit, canopy and root architecture, plant population density, and water and nutrient usage efficiency are the primary elements that influence intercropping. Crop productivity and efficiency per unit area of land have been measured using a variety of competitive indicators. Land Equivalent Ratio, Area Time Equivalent Ratio, Aggressivity, Relative crowding coefficient, Competitive ratio, Actual yield loss are the most frequent competitive index approaches. Generally intercropping should be widely used in developing nations to provide agricultural sustainability, food security, and nutrition.

## References

1. Abera T, Feyissa D, Yusuf H. Effects of inorganic and organic fertilizers on grain yield of maize-climbing bean intercropping and soil fertility in western Oromiya, Ethiopia. Tropentag 2005 Stuttgart-Hohenheim, October 11-13, 2005. Conference on International Agricultural Research for Development. 2005.
2. Addisu Getahun, Seltene Abady. Effect of Maize (*Zea mays* L.) on Bean (*Phaseolus vulgaris* L.) Yield and its Components in Maize-Bean Intercropping. International Journal of Science and Research (IJSR): ISSN: 2319-7064. 2014.
3. Addo-Quaye AA, Darkwa AA, Ocloo GK. Yield and productivity of component crops in a maize-soybean intercropping system as affected by time of planting and spatial arrangement. Journal of Agricultural and Biological Science. 2011;6(9):50-57.
4. Adu-Gyamfi JJ, Myaka FA, Sakala WD, Odgaard R, Vesterager JM, Høgh-Jensen H. Biological nitrogen fixation and nitrogen and phosphorus budgets in farmer-managed intercrops of maize-pigeonpea in semi-arid Southern and Eastern Africa. Plant Soil. 2007;295:127-136.
5. Alemayehu D, Shumi D, Afeta T. Effect of Variety and Time of Intercropping of Common Bean (*Phaseolus vulgaris* L.) With Maize (*Zea mays* L.) on Yield Components and Yields of Associated Crops and Productivity of the System at Mid-Land of Guji, Southern Ethiopia. Adv Crop Sci Tech. 2018;6:324. DOI: 10.4172/2329-8863.1000324
6. Allen JR, Obura RK. Yield of corn and soybean in intercropping soybeans. Agron. J. 1983;75:1005-1009.
7. Anders MM, Potdar MV, Francis CA. The significance of Intercropping in cropping systems. In: Ito, O., Johansen C, Adu-Gyamfi JJ, Katayama K, Kumar JVD, Rao K, Rego TJ. (Eds.). Dynamics of roots and nitrogen in cropping systems of the semi-arid tropics. Japan International Research Center for Agricultural Sciences. International Agricultural Series No. 3 Ohwashi, Tsukuba, Ibaraki 305, Japan. 1996.
8. Balde AB, Scope LE, Affholder F, Corbeels M, DaSilva FAM, Xavier JHV, *et al.* Agronomic performance of no-tillage relay intercropping with maize under smallholder conditions in Central Brazil. Field Crop Res. 2011;124:240-251. [Cross Ref]
9. Banik P, Sasmal T, Ghosal PK, Bagchi DK. Evaluation of Mustard (*Brassica campestris* var. Toria) and legume intercropping under 1:1 and 2:1 row-replacement series systems. J Agron. Crop Sci. 2000;185:9-14.
10. Banik P, Midya A, Sarkar BK, Ghose SS. Wheat and chickpea intercropping systems in an additive series experiment: advantages and weed smothering. Eur. J Agron. 2006;24:325-332.
11. Chalka MK, Nepalia V. Nutrient uptake appraisal of maize intercropped with legumes and associated weeds under the influenced of weed control. Indian Journal of Agricultural Research. 2006;40(2):86-91.
12. Dereje G, Adisu T, Mengesha M, Bogale T. The Influence of Intercropping Sorghum with Legumes for Management and Control of Striga in Sorghum at Assosa Zone, Benshangul Gumuz Region, Western Ethiopia, East Africa. Adv Crop Sci Tech. 2016;4:238. DOI: 10.4172/2329-8863.1000238.
13. Dahmardeh M, Ghanbari A, Syahsar BA, Ramrodi M. The role of intercropping maize (*Zea mays* L.) and cowpea (*Vigna unguiculata* L.) on yield and soil chemical properties. African Journal of Agricultural Research. 2010;5(8):631-636. Available online at <http://www.academicjournals.org/AJAR>. ISSN 1991-637X © 2010 Academic Journals.
14. Eskandari H, Ghanbari A, Javanmard A. Intercropping of cereals and legumes for forage production. Not. Sci. Biol. 2009, 2010;1:7-13. [Cross Ref]
15. Francis CA. Introduction: distribution and importance of multiple cropping. Pages 1-20 in Multiple cropping systems (Francis, C.A., ed.). Macmillan Publishing Company, New York, USA. 1986, 1-20.
16. Fujita K, Ofosu-Budu KG. Significance of Intercropping in Cropping Systems. 1996, 19-40.
17. Gebremichael A, Bekele B, Tadesse B. Evaluation of the effect of sorghum-legume intercropping and its residual effect on yield of sorghum in yekiworeda, sheka zone, Ethiopia. Int. J. Agril. Res. Innov. Tech. 2019;9(2):62-66. DOI: 10.3329/ijarit.v9i2.45412.
18. Gulwa U, Mgujulwa N, Beyene ST. Effect of Grass-legume Intercropping on Dry Matter Yield and Nutritive Value of Pastures in the Eastern Cape Province, South Africa. Univ. J Agric. Res. 2017;5:355-362. [Cross Ref]
19. Hailu Gebru. A Review on the Comparative Advantages of Intercropping to Mono-Cropping System. Journal of Biology, Agriculture and Healthcare. ISSN 2224-3208. 2015, 5(9).
20. Harris DR. Traditional systems of plant food production and the origins of agriculture in West Africa. Pages 311-346 in Origins of african plant domestication (Harlan, J.R., De Wet, J.M.J., and Stemler, A.B.L., eds.). Mouton, The Hague. 1976.



21. Jaya IKD, Bell CJ, Sale PW. Modification of within-microclimate in maize for intercropping in the lowland tropics. 123-133. Proceedings of the Australian Agronomy Conference. Australia Society of Agronomy, Australia. 2001.
22. Keating B, Carberry P. Resource capture and use in intercropping: Solar radiation. *Field Crops Research*. 1993;34:273-301.
23. Li CJ, Hoffland E, Kuyper TW, Yu Y, Zhang C, Li H, *et al.* Syndromes of production in intercropping impact yield gains. *Nature Plants*. 2020a;6:653-660. <https://doi.org/10.1038/s41477-020-0680-9>
24. Li CJ, Hoffland E, Kuyper TW, Yu Y, Li H, Zhang C, *et al.* Yield gain, complementarity and competitive dominance in intercropping in China: A meta-analysis of drivers of yield gain using additive partitioning. *Eur J Agron* 2020b;113:125987. <https://doi.org/10.1016/j.eja.2019.125987>
25. Martin-Guay M, Paquette A, Dupras J, Rivest D. The new green revolution: sustainable intensification of agriculture by intercropping. *Sci Total Environ*. 2018;615:767-772.
26. Mashingaidze AB. Improving weed management and crop productivity in maize systems in Zimbabwe. Ph. D thesis, Wageningen University. 2004, 207.
27. Moris RA, Garrity D1'. Resource capture and utilization in intercropping: non-nitrogen nutrients. *Field Crops Research*. 1993a, 34319-334.
28. Mucheru-Muna M, Pypers P, Mugendi D, Kung'u J, Mugwe J, Merckx R, Vanlauwe B. Staggered maize-legume intercrop arrangement robustly increases crop yields and economic returns in the highlands of Central Kenya. *Field Crops Research*. 2010;115:132-139.
29. Musambasi D, Chivinge OA, Mariga IK. Intercropping Maize with Grain Legumes for Striga Control in Zimbabwe. *African Crop Science Journal*. 2002;10(2):163-171.
30. Ofori F, Stern WR. Cereal-legume intercropping systems. *Advances in Agronomy*. 1987;40:41-90.
31. Osman AN, Ræbild A, Christiansen JL, Bayala J. Performance of cowpea (*Vigna unguiculata*) and Pearl Millet (*Pennisetum glaucum*) Intercropped under Parkia. 2011.
32. Plucknett DL, Smith NJH. Historical perspectives on multiple cropping. Pages 20-39 in *Multiple cropping systems* (Francis C.A., ed.). Macmillan Publishing Company. 1986, 20-39.
33. Sangina N, Woomer PL. Integrated soil fertility management in Africa: Principles, Practices and Development Process. (Eds.). *Tropical Soil Biology and Fertility Institute of the International Centre for Tropical Agriculture*. Nairobi. 2009, 263.
34. Sarkar RK, Shit D, Chakraborty A. Yield and economics of pigeonpea based intercropping system on rainfed upland of Chotanagpur plateau. *Indian Journal of Agronomy*. 1995;40:30-34.
35. Seran TH, Brintha I. Review on maize based intercropping. *Journal of Agronomy*. 2010;9(3):135-145.
36. Synadon RW, Harris PM. Interactions below ground the use of nutrients and water. Proceedings of the International Work Shop on Intercropping (IWSI'79), ICRISAT, India. 1979, 181-201.
37. Ananthi T, Mohamed Amanullah M, Abdel Rahman Mohammad Said Al-Tawaha. A Review on Maize- Legume Intercropping For Enhancing The Productivity And Soil Fertility For Sustainable Agriculture In India. *Advances in Environmental Biology*. 2017;11(5):49-63.
38. Tamiru Hirpa, Heluf Gebrekidan, Kindie Tesfaye, Asfaw Hailemariam. Production efficiency of maize-based cropping system as affected by intercropping date of companion legume crops. *Ethiop. J Biol. Sci*. 2013;12(1):51-65.
39. Tana T, Mulatu E. Evaluation of sorghum, maize and common bean cropping system in east Hararghe, Eastern Ethiopia. *Ethiopian Journal of Agricultural Science*. 2000;17:33-46.
40. Tenaw Workayehu, Legesse Hidoto, Gobeze Loha. Grain Yield and Economic Benefit of Intercropping Barley and Faba Bean in the Highlands of Southern Ethiopia. *East African Journal of Sciences*. 2016;10(2):103-110.
41. Undie UL, Uwah DF, Attoe E. Effect of intercropping and crop arrangement on yield and productivity of late season maize/soybean mixtures in the humid environment of south southern Nigeria. *J Agric. Res*. 2012;4:37. [Cross Ref]
42. Varma D, Meena RS, Kumar S. Response of mungbean to fertility and lime levels under soil acidity in an alley cropping system in Vindhyan region. *Indian Int. J Chem. Stud*. 2017;5:384-389.
43. Vesterager JM, Nielsen NE, Høgh-Jensen H. Effect of cropping history and phosphorous source on yield and nitrogen fixation in sole and intercropped cowpea-maize systems. *Nutrient Cycling in Agroecosystems*. 2008;80:61-73.
44. Von Cossel M, Wagner M, Lask J, Magenau E, Bauerle A, Von Cossel V, *et al.* Prospects of Bioenergy Cropping Systems for A More Social-Ecologically Sound Bioeconomy. *Agronomy*. 2019;9:605. [Cross Ref]
45. Westphal E. Pulses in Ethiopia, Their Taxonomy and Agricultural Significance; Centre for Agricultural Publishing and Documentation: Wageningen, The Netherlands. 1974, 275.
46. Willey RW. Intercropping- its importance and research needs. Part 1: Competition and yield advantages. *Field Crop Research*. 1979;32:1-10.
47. Willey RW, Rao MR, Natarajan M. Traditional cropping systems with pigeonpea and their improvement. In: Nene, Y. L. and Kumble, V. ICRISAT. 1981. Proceedings of the International Workshop on Pigeonpeas 1:15-19 December 1980, Patancheru, A.P., India. 1980.
48. Willey RW. Resource use in intercropping systems. *Journal of Agriculture and Water Management*. 1990;17:215-231.
49. Yang F, Wang XC, Liao DP, Lu FZ, Gao RC, Liu WG, *et al.* Yield response to different planting geometries in maize-soybean relay strip intercropping systems. *Agron. J*. 2015;107:296-304. [Cross Ref]
50. Yu Y, Stomph TJ, Makowski D, Van der Werf W. Temporal niche differentiation increases the land equivalent ratio of annual intercrops: A meta-analysis. *Field Crops Res*. 2015;184:133-144.